
NOTE: The horizontal line across the page indicates the maximum allowed length of the abstract including footnotes. If your abstract is close to or exceeds the length limit, it may appear on page 2 of this document.

Time required: 15 min

Session Categories (Keywords) by Relevance: Small molecules — Perturbations, resonances, *-Teller — Electronic structure, potential energy surfaces — Optical/UV/X-ray — Fundamental interest

Mini-Symposia Requested: None — None

Competing for Rao Prize? No

Competing for Miller Prize? No

DIRECT POTENTIAL FIT FOR THE $X^1\Sigma$ STATE OF F_2 : PERTURBATION OF THE HIGHEST OBSERVED $v=22$ VIBRATIONAL LEVEL

ROBERT W FIELD, *Department of Chemistry, MIT, Cambridge, MA, USA*; JOHN COXON, *Department of Chemistry, Dalhousie University, Halifax, NS, Canada*; PHOTOS HAJIGEORGIOU, *Centre for Primary Care and Population Health, School of Medicine, University of Nicosia, Nicosia, Cyprus*.

The high-resolution vacuum-uv spectrographic data¹ for the C - X emission and C,(D,E),H,h,I - X($v = 0$) absorption transitions of F_2 , in combination with pure rotation² and vibration-rotation³ Raman data, have been employed in a least-squares analysis. Attention was given to the extensive blending in the absorption data and to account for plate-to-plate shifts in the emission data. The C - X data, with an estimated uncertainty of 0.05 cm^{-1} , sample X state vibrational levels $v = 1 - 22$, for which the potential energy function was fitted using the extended-MLR model⁴. 3549 line positions in the weighted fit provided estimates of 1303 term values of excited electronic states and 17 parameters for the ground state. The highest observed $v = 22$ level of the ground state, which lies only 114 cm^{-1} below the $F(2P_{3/2}) + F(2P_{3/2})$ dissociation limit, is found to be perturbed; all rotational levels ($J = 0 - 19$) lie at energies $5 - 13 \text{ cm}^{-1}$ below their expected positions. A deperturbation model was employed within the direct potential fit; in this novel approach, the eigenvalue of each J-level in $v = 22$ was determined from a 2×2 matrix, with the diagonal level of the perturbing state represented by $E_p + B_p J(J+1)$, and the off-diagonal element by $a + b(J + 1/2)$. However, the b-parameter was indeterminate; a successful fit of the entire data set with inclusion of the deperturbation model for $v = 22$ provided the estimates $E_p = -70.5(3.7) \text{ cm}^{-1}$, $B_p = 0.226(6) \text{ cm}^{-1}$, $a = 16.2(8) \text{ cm}^{-1}$ and $R_e = 1.412555(4) \text{ \AA}$. There is much interest in an identification of the perturbing state. The results indicate a J-independent spin-orbit interaction with a weakly-bound perturbing state ($R_e = 2.8 \text{ \AA}$), lying $40 - 50 \text{ cm}^{-1}$ above $v = 22$. The absence of a J-dependent $b(J + 1/2)$ Coriolis interaction implies a perturber with 0_g^+ symmetry. A plausible candidate is the $a'(0_g^+)$ state which dissociates to the same atomic limit and which is repulsive at short-R.

1. E.A. Colbourn, M. Dagenais, A.E. Douglas, J.W. Raymonda, *Can. J. Phys.* 54 (13) (1976) 1343-1359. 2. H.G.M. Edwards, E.A.M. Good, D.A. Long, *J. Chem. Soc. Faraday Trans.* 272 (1976) 984-987. 3. R.Z. Martinez, D. Bermejo, J. Santos, P. Cancio, *J. Mol. Spectrosc.* 168 (1994) 343-349. 4. R.J. Le Roy, N.S. Dattani, J.A. Coxon, A.J. Ross, P. Crozet, C. Linton, *J. Chem. Phys.* 131 (2009) 204309.

NOTE: The horizontal line across the page indicates the maximum allowed length of the abstract including footnotes. If your abstract is close to or exceeds the length limit, it may appear on page 2 of this document.

Time required: 15 min

Session Categories (Keywords) by Relevance: Small molecules — Perturbations, resonances, *-Teller — Electronic structure, potential energy surfaces — Optical/UV/X-ray — Fundamental interest

Mini-Symposia Requested: None — None

Competing for Rao Prize? No

Competing for Miller Prize? No