

ULTRAVIOLET PHOTOABSORPTION OF SO ISOTOPOLOGUES AND THE $B^3\Sigma^-$ AND $C^3\Pi$ STATES

ALAN HEAYS^a, *School of Earth and Space Exploration, Arizona State University, Tempe, AZ, USA*; GLENN STARK, *Department of Physics, Wellesley College, Wellesley, MA, USA*; JAMES R LYONS, *School of Earth and Space Exploration, Arizona State University, Tempe, AZ, USA*; NELSON DE OLIVEIRA, *DESIRS Beamline, Synchrotron SOLEIL, Saint Aubin, France*; BRENTON R LEWIS, STEPHEN T GIBSON, *Research School of Physics and Engineering, Australian National University, Canberra, ACT, Australia*.

The sulphur-monoxide $B^3\Sigma^-(v \geq 4)$ levels are known to be strongly affected by vibrationally-dependent predissociation and local energy perturbations (Liu et al. 2006 JMS 238:213). The isotope-dependence of this predissociation and the SO photodissociation cross section is a candidate atmospheric-source for explaining the anomalous $^{32}\text{S}/^{33}\text{S}/^{34}\text{S}/^{36}\text{S}$ isotopic fractionation found in 2.5Ga old sedimentary material (Ono 2017 Annu. Rev. Earth Pl. Sc. 45:301).

We have recorded new photoabsorption spectra between 195 and 230 nm to determine spectroscopic constants, predissociation linewidths, and transition strengths for the excited $B^3\Sigma^-(v = 4-17)$ levels of $^{32}\text{S}^{16}\text{O}$, $^{33}\text{S}^{16}\text{O}$, and $^{34}\text{S}^{16}\text{O}$. The $C^3\Pi$ state is also observed and perturbs $B^3\Sigma^-(v = 4-17)$ through spin-orbit interaction. $B^3\Sigma^-$ and $C^3\Pi$ potential-energy curves, electronic transition moments, and a global spin-orbit interaction are deduced from the new data so that it may be extrapolated to the rare $^{36}\text{S}^{16}\text{O}$ isotopologue.

We use the new cross sections to explore the potential for isotope-dependent photodissociation of SO in the ancient-Earth atmosphere due to structured solar UV radiation and atmospheric opacity.

^aNASA Astrobiology Institute, NASA Ames Research Center, Moffett Field, California