



I L L I N O I S

UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN

**PRODUCTION NOTE**

University of Illinois at  
Urbana-Champaign Library  
Large-scale Digitization Project, 2007.

**This page is intentionally blank.**

DOC  
ILNHS  
FAU  
1985(2)

Natural History Survey  
Library

# ILLINOIS NATURAL HISTORY SURVEY

SURVEY OF KANKAKEE RIVER FISHES AND MUSSELS AT WASHINGTON AVENUE BRIDGE,  
FAU 6198, U. S. ROUTES 45 AND 52,  
CITY OF KANKAKEE, KANKAKEE COUNTY, ILLINOIS.  
IDOT JOB NUMBER P-93-055-83; STRUCTURE NUMBER 046-0049

Final Report

4 April 1985




## Section of Faunistic Surveys and Insect Identification Technical Report

by

Mark J. Wetzel  
Jeanine M. Kasprowicz  
Kevin S. Cummings  
Warren U. Brigham  
Liane Suloway

Prepared for

Illinois Department of Transportation  
Bureau of Location and Environment  
2300 South Dirksen Parkway  
Springfield, Il 62764



Section of Faunistic Surveys and Insect Identification  
Technical Report 1985(2)



## TABLE OF CONTENTS

	PAGE
INTRODUCTION .....	1
DESCRIPTION OF STUDY AREA .....	1
FISH SURVEY .....	3
Methods .....	3
Results .....	3
Discussion .....	3
LIFE HISTORY ACCOUNTS OF THREATENED ILLINOIS FISHES FOUND IN THE KANKAKEE RIVER SYSTEM .....	8
MUSSEL SURVEY .....	10
Methods .....	10
Results .....	11
Discussion .....	11
PRESENT STATUS OF MUSSELS IN THE KANKAKEE RIVER .....	16
ACKNOWLEDGEMENTS .....	17
LITERATURE CITED .....	17

## INTRODUCTION

The Kankakee River has long been recognized as one of the highest-quality streams in Illinois (Ivens *et al.* 1981). It supports a large and diverse mussel population which has been studied intermittently for nearly 100 years (Baker 1906; Wilson and Clark 1912; Parmalee 1967; Brice and Lewis 1979; Lewis and Brice 1980; Page *et al.* [1979]; Brigham, Suloway, and Page 1981; Suloway 1981; Brigham *et al.* 1984, Kasprowicz *et al.* 1984; and Wetzel, Kasprowicz, and Brigham 1984).

In addition, one species of fish known to occur in the Kankakee River - the blacknose shiner (*Notropis heterolepis* Eigenmann and Eigenmann) - is a state threatened species (Smith and Page 1981). The pallid shiner (*Notropis amnis* Hubbs and Greene), a fish previously thought to be extirpated from Illinois, but recently discovered in the Kankakee River (Skelly and Sule 1983), has been proposed for listing as a state threatened species (Sweet 1984).

The Illinois Department of Transportation (IDOT), Bureau of Location and Environment, is required to comply with Section 404(b)1 of the Clean Water Act when projects have the potential to impact on aquatic environments. In addition, IDOT also is required by Section 7 of the Endangered Species Act of 1973, as amended, to conduct a biological inventory for the purpose of identifying federally listed endangered and threatened species likely to be affected by its highway construction projects. IDOT also wishes to consider in a similar fashion those species listed as endangered or threatened in Illinois by the state Endangered Species Protection Board. In many cases, the general distribution patterns of these species are known, but specific records from the project area are not available. Usually a field reconnaissance and review of appropriate publications and museum records are necessary to determine whether or not a species actually occurs or is thought likely to occur in a project area.

In conjunction with the proposed IDOT project FAU 6198, in which the existing Washington Avenue bridge over the Kankakee River in the City of Kankakee, Kankakee County, Illinois will be removed and replaced, the Illinois Natural History Survey conducted such an inventory to determine the actual or likely occurrence of mussel and fish species within the project area. Particular attention was given to federal and state endangered and threatened species, which, based upon past records and information about habitat preferences, were known or thought likely to occur in the Kankakee River.

## DESCRIPTION OF STUDY AREA

The Washington Avenue bridge is located in the City of Kankakee, 90 m downstream of the Kankakee River dam and hydroelectric plant, and carries U. S. routes 45 and 52 across the river (Figure 1). The legal location of the study area, taken from a U. S. Geological Survey topographic quadrangle map, is as follows:

IL, Kankakee Co., Kankakee River, in Kankakee. T.30 N, R.13 W,  
NW/4 & SW/4, Sec. 5. U.T.M.: 4551500m N, 427100m E, Zone 16.  
Kankakee, Ill. (7.5' series, 1964 ed, 1973PR) USGS quad.




Figure 1. Location of study site discussed in *SURVEY OF KANKAKEE RIVER FISHES AND MUSSELS AT WASHINGTON AVENUE BRIDGE, FAU 0198, U. S. ROUTES 45 AND 52, CITY OF KANKAKEE, KANKAKEE COUNTY, ILLINOIS*. IDOT JOB NUMBER P-93-055-83; STRUCTURE NUMBER 040-0049 (from Kankakee, Ill. 7.5' quadrangle).

## FISH SURVEY

Methods

Fishes were sampled using a 4'x10'x1/8" common minnow seine. Fifteen seine hauls were taken at the study site (Figure 1). Habitats sampled included pools, riffles, and raceways. Sand and silt substrates predominated in pools and backwater areas while gravel, rocks, and bedrock were most common in raceways and riffles. Fish collections were made on 21 September 1984 under the supervision of INHS ichthyologist Kevin S. Cummings, with the assistance of INHS zoologists Mark J. Wetzel and Jeffrey T. Irish.

Results

A total of 12 species was collected from this site, representing 6 families and eight genera (Table 1). Minnows accounted for 50% of the species taken, with spotfin shiners (*Notropis spilopterus* (Cope)), bluntnose minnows (*Pimephales notatus* (Rafinesque)), and sand shiners (*Notropis stramineus* (Cope)) accounting for 140 of the 169 specimens collected.

A total of 86 species of fishes has been collected in the Kankakee River in recent times (Page *et al.* [1979]). Because of habitat preference, some fishes are only found in the headwaters while others only downstream. The combination of low water level, lack of available habitat, and selective collecting apparatus probably accounted for the smaller number of species collected in 1984.

Discussion

Of all the fishes known to occur in the Kankakee River (Table 2), only one, the blacknose shiner (*Notropis heterolepis* Eigenmann & Eigenmann), is listed as threatened in the State of Illinois (Smith and Page 1981). However, the blacknose shiner is restricted primarily to tributaries of the Kankakee River and is seldom found in the river mainstream.

Another species known to occur in the Kankakee River and of special concern is the pallid shiner (*Notropis amnis* Hubbs & Greene). This species, once thought to have been extirpated from Illinois, was collected in 1978, 1979, 1981, and 1982 from the Kankakee River near Custer Park (Skelly and Sule 1983). Because of its presence in the Kankakee River, the pallid shiner is now being considered for the Illinois list of threatened species (Sweet 1984). This species lives in quiet silt- and sand-bottomed pools of large rivers. The population remaining in the Kankakee River persists because of high water quality and suitable habitat.

Life history accounts of the blacknose shiner and the pallid shiner are included below.

Table 1. List of fishes collected 21 September 1984 from the Kankakee River at the Washington Avenue bridge, City of Kankakee, Kankakee County, Illinois.

Family Cyprinidae - carps and minnows	Number collected
<i>Nocomis biguttatus</i> (Kirtland)	1
<i>Notropis chrysocephalus</i> (Rafinesque)	3
<i>Notropis lutrensis</i> (Baird & Girard)	2
<i>Notropis spilopterus</i> (Cope)	117
<i>Notropis stramineus</i> (Cope)	7
<i>Pimephales notatus</i> (Rafinesque)	16
Family Ictaluridae - bullhead catfishes	
<i>Noturus flavus</i> Rafinesque	2
Family Cyprinodontidae - killifishes	
<i>Fundulus notatus</i> (Rafinesque)	5
Family Atherinidae - silversides	
<i>Labidesthes sicculus</i> (Cope)	1
Family Centrarchidae - sunfishes	
<i>Lepomis macrochirus</i> Rafinesque	10
Family Percidae - perches	
<i>Etheostoma nigrum</i> Rafinesque	2
<i>Etheostoma zonale</i> (Cope)	3
Total	169

Table 2. List of fishes known to occur in the Kankakee River, Illinois.  
 (Sources: Illinois Department of Conservation, Illinois Natural  
 History Survey, Page *et al.* [1979], Brigham *et al.* 1981).

FAMILY	SCIENTIFIC NAME	COMMON NAME
PETROMYZONTIDAE lampreys	<i>Ichthyomyzon fossor</i> <i>I. unicuspis</i> <i>Lampretra appendix</i>	northern brook lamprey silver lamprey American brook lamprey
LEPISOSTEIDAE gars	<i>Lepisosteus osseus</i>	longnose gar
AMIIDAE bowfins	<i>Amia calva</i>	bowfin
ANGUILLIDAE freshwater eels	<i>Anguilla rostrata</i>	American eel
HIODONTIDAE mooneyes	<i>Hiodon alosoides</i>	goldeye
CLUPEIDAE herrings	<i>Alosa chrysochloris</i> <i>Dorosoma cepedianum</i>	skipjack herring gizzard shad
SALMONIDAE trouts	<i>Salmo gairdneri</i>	rainbow trout
UMBRIDAE mudminnows	<i>Umbra limi</i>	central mudminnow
ESOCIDAE pikes	<i>Esox americanus</i> <i>E. lucius</i> <i>E. masquinongy</i>	grass pickerel northern pike muskellunge
CYPRINIDAE carps & minnows	<i>Campostoma anomalum</i> <i>Carassius auratus</i> <i>Cyprinus carpio</i> <i>Ericymba buccata</i> <i>Nocomis biguttatus</i> <i>Notemigonus crysoleucas</i> <i>Notropis amnis</i> <i>N. atherinoides</i> <i>N. buechanani</i> <i>N. chalybaeus</i> <i>N. chrysocephalus</i> <i>N. cornutus</i> <i>N. dorsalis</i> <i>N. emiliae</i> <i>N. heterolepis</i>	central stoneroller goldfish common carp silverjaw minnow hornyhead chub golden shiner pallid shiner emerald shiner ghost shiner ironcolor shiner striped shiner common shiner bigmouth shiner pugnose minnow blacknose shiner

Table 2. Continued.

FAMILY	SCIENTIFIC NAME	COMMON NAME
CYPRINIDAE (cont'd)	<i>N. lutrensis</i>	red shiner
	<i>N. rubellus</i>	rosyface shiner
	<i>N. spilopterus</i>	spotfin shiner
	<i>N. stramineus</i>	sand shiner
	<i>N. texanus</i>	weed shiner
	<i>N. umbratilis</i>	redfin shiner
	<i>N. volucellus</i>	mimic shiner
	<i>Phenacobius mirabilis</i>	suckermouth minnow
	<i>Pimephales notatus</i>	bluntnose minnow
	<i>P. promelas</i>	fathead minnow
	<i>P. vigilax</i>	bullhead minnow
	<i>Semotilus atromaculatus</i>	creek chub
	<i>Cyprinus carpio</i> x <i>Carassius auratus</i>	hybrid
CATOSTOMIDAE suckers	<i>Carpiodes carpio</i>	river carpsucker
	<i>C. cyprinus</i>	quillback
	<i>Catostomus commersoni</i>	white sucker
	<i>Erimyzon sucetta</i>	lake chubsucker
	<i>Hypentelium nigricans</i>	northern hog sucker
	<i>Ictiobus bubalus</i>	smallmouth buffalo
	<i>I. cyprinellus</i>	bigmouth buffalo
	<i>I. niger</i>	black buffalo
	<i>Minytrema melanops</i>	spotted sucker
	<i>Moxostoma anisurum</i>	silver redhorse
	<i>M. carinatum</i>	river redhorse
	<i>M. erythrurum</i> <i>M. macrolepidotum</i>	golden redhorse shorthead redhorse
ICTALURIDAE bullhead catfishes	<i>Ictalurus melas</i>	black bullhead
	<i>I. natalis</i>	yellow bullhead
	<i>I. punctatus</i>	channel catfish
	<i>Noturus flavus</i>	stonecat
	<i>Pylodictis olivaris</i>	flathead catfish
CYPRINODONTIDAE killifishes	<i>Fundulus notti</i>	starhead topminnow
	<i>F. notatus</i>	blackstripe topminnow
ATHERINIDAE silversides	<i>Labidesthes sicculus</i>	brook silversides
APHREDODERIDAE pirate perches	<i>Aphredoderus sayanus</i>	pirate perch

Table 2. Concluded.

FAMILY	SCIENTIFIC NAME	COMMON NAME
CENTRARCHIDAE sunfishes	<i>Ambloplites rupestris</i>	rock bass
	<i>Lepomis cyanellus</i>	green sunfish
	<i>L. gibbosus</i>	pumpkinseed
	<i>L. gulosus</i>	warmouth
	<i>L. humilis</i>	orangespotted sunfish
	<i>L. macrochirus</i>	bluegill
	<i>L. megalotis</i>	longear sunfish
	<i>L. microlophus</i>	redeer sunfish
	<i>Micropterus dolomieu</i>	smallmouth bass
	<i>M. salmoides</i>	largemouth bass
	<i>Pomoxis annularis</i>	white crappie
	<i>P. nigromaculatus</i>	black crappie
	<i>Lepomis megalotis</i> x ?	hybrid
PERCIDAE perches	<i>Etheostoma caeruleum</i>	rainbow darter
	<i>E. chlorosomum</i>	bluntnose darter
	<i>E. flabellare</i>	fantail darter
	<i>E. microperca</i>	least darter
	<i>E. nigrum</i>	johnny darter
	<i>E. zonale</i>	banded darter
	<i>Perca flavescens</i>	yellow perch
	<i>Percina caprodes</i>	logperch
	<i>P. maculata</i>	blackside darter
	<i>P. phoxocephala</i>	slenderhead darter
<i>Stizostedion vitreum</i>	walleye	

LIFE HISTORY ACCOUNTS OF THREATENED ILLINOIS FISHES  
FOUND IN THE KANKAKEE RIVER SYSTEM*Notropis heterolepis* Eigenmann & Eigenmann - Blacknose shiner

The blacknose shiner is a northern minnow, ranging from Saskatchewan to Nova Scotia, but persisting in disjunct relict populations as far south as Illinois, Indiana, Missouri, and Ohio. Cross (1967) suggested that this species was extirpated recently from Kansas. Formerly, this species was found throughout the northern two-thirds of Illinois, and in Clear Creek, Union County (Forbes and Richardson 1920). Presently, however, the blacknose shiner is known only from glacial lakes and tributaries of the Fox River in Lake and McHenry counties, the Kankakee River drainage in Iroquois, Kankakee, and Will counties, from Fairfield Ditch, Green River drainage in Bureau County, the Kishwaukee River in Boone and McHenry counties, and from Coon Creek in Whiteside County (Smith 1979; Smith and Page 1981).

No comprehensive systematic review of *Notropis heterolepis* has been done (Gilbert 1980). Presently, only the nominate subspecies, *N. heterolepis heterolepis* is known to occur in Illinois (see Scott and Crossman 1973, and Smith 1979 for additional information).

The blacknose shiner is known to occur in cool, clear, densely vegetated waters (Gilbert 1980; Smith 1979; Becker 1983), including lakes and streams (Smith and Page 1981).

Although no detailed information on the food habits of the blacknose shiner have been published, insect larvae and entomostracans have been found to form a large part of the diet (Cahn 1927; Becker 1983). In Minnesota, algae, sponges, and larval and adult insects were the principal food items found (Nurnberger 1928). However, the ventral position of the mouth and short intestine suggests that this species may be a carnivorous benthic feeder (Becker 1983).

While limiting factors of reproduction are unknown, the blacknose shiner appears to be extremely intolerant of siltation and turbidity (Trautman 1957; Scott and Crossman 1973; Smith 1979; Smith and Page 1981; Becker 1983). Breeding season in Illinois is from June to August (Forbes and Richardson 1920; Smith 1979). Spawning behavior and habits are unknown, but it has been suggested that spawning occurs over sandy areas (Scott and Crossman 1973; Becker 1983). Lopinot and Smith (1973) reported that this species reaches sexual maturity and spawns at 1 year of age.

Reason for threatened status in Illinois: The disappearance of the blacknose shiner in Illinois has been among the most dramatic of any fish (Smith and Page 1981, and elsewhere). The blacknose shiner is no longer present in many localities in which it formerly occurred (Trautman 1957; Scott and Crossman 1973; Gilbert 1980; Smith 1979; Smith and Page 1981; Becker 1983). Increased siltation, turbidity, and the disappearance of aquatic vegetation are probably the major factors causing its decline in Illinois (Smith 1979). These factors have been shown to be responsible for the decline of this species in Ohio (Trautman 1957), and in Missouri (Pflieger 1975).

*Notropis amnis* Hubbs & Green - Pallid shiner

The pallid shiner, *Notropis amnis* Hubbs and Greene, is one of the rarest and least known American fishes (Smith 1979). Its range extends from the Mississippi River in Minnesota and Wisconsin south to the Amite River in Louisiana and west to the Guadalupe River in Texas (Clemmer 1980). This fish is rare in the northern half of its range; since 1970 most specimens have been taken from backwaters of the Mississippi River in Wisconsin, primarily between the Wisconsin-Illinois border and Prairie du Chien (Don Fago, Wisconsin Department of Natural Resources, pers. comm. to Skelly and Sule 1983). This species has shown the most marked decline of any fish in Missouri and may have been eliminated from that state (Pfleiger 1975). It was thought to be extirpated from Illinois until its recent collection in the Kankakee River. Since 1978, Illinois Natural History Survey biologists have collected 18 specimens, 15 in August collections and 3 in November. Prior to these collections only 22 specimens of *N. amnis* were considered Illinois records, the last collected in 1963 (Skelly and Sule 1983).

*Notropis amnis* from the Kankakee River was found in areas of slight current, no measurable current, and backflow. Although turbidities during collecting periods were low, the river is subject to dramatic increases in turbidity (as high as 300 NTU) during storm periods. All *N. amnis* were collected in seizable areas at water depths no greater than 1.5 m. Water temperatures varied seasonally, with specimens collected in water 6.0°C to 29.5°C. These quantitative data supplement earlier observations, such as turbidity ranging from "clear" to "very muddy" (Hubbs 1951), and that *N. amnis* is somewhat tolerant of turbidity (Clay 1975). Pflieger (1975) stated that this species is intolerant of excessive siltation and turbidity. Several studies have shown that *N. amnis* resides in slowly moving waters (Hubbs 1951; Miller and Robison 1973; Clay 1975; and Pflieger 1975). When collected from the Mississippi River, it has been collected in flowing water over sand bars (Harlan and Speaker 1969; Eddy and Underhill 1974).

Most of the specimens of *Notropis amnis* collected by Skelly and Sule (1983), were taken from an area between 200 and 750 m downstream of the confluence of Horse Creek with the Kankakee River near Custer Park. Unlike other stations within the study area that were periodically scoured by currents, this area remained a slack-water, depositional area despite fluctuating levels of river discharge. Year-round slow current and depositional substrate appear to be the preferred habitat of *N. amnis*, since no specimens were collected by Skelly and Sule in any other habitat type despite extensive sampling throughout the monitoring area.

Good water quality throughout the study area, as indicated by the analysis of 54 water quality parameters (Skelly and Sule 1980), and a variety of habitats are important to the maintenance of the diverse Kankakee River assemblage of fishes, totaling at least 86 species. The Kankakee River in Illinois is noted for its good water quality, diverse aquatic life, and scenic beauty (Ivens *et al.* 1981).

*Notropis amnis* seldom enters the mouths of tributary streams (Hubbs 1951; Eddy and Underhill 1974). The lentic characteristics within the mouth of Horse Creek, a third order stream (Horton-Strahler system) entering the Kankakee River within the study area of Skelly and Sule (1983), were quite similar to

those found between its mouth and 750 m downstream of its confluence, yet no pallid shiners were collected from that creek. Hubbs (1951) suggested that *N. ammis* in its northern range is confined to the large lowland rivers, possibly because large rivers are warmer than small to medium-sized streams. Small streams like Horse Creek are also subject to sudden increases in current that may be unacceptable to *N. ammis*.

Thirty-six species representing 8 families of fishes were collected by Skelly and Sule (1983) at stations where *N. ammis* was collected. Species and abundance of fishes collected varied by date and station. However, the most common associations and those found during each sampling period included *Notropis spiloferus* and *Pimephales notatus*. *Pimephales vigilax* and *Notropis stramineus* were abundant during several collections, and *Lepomis humilis* was occasionally common. *Notropis spiloferus* is the only member of this group that is considered characteristic of high-gradient streams. The other species are characteristic of quiet pools and backwaters of medium to moderately large streams. Both *N. spiloferus* and *L. humilis* are somewhat tolerant of turbid conditions (Pflieger 1975).

*Notropis ammis* has not been officially listed as endangered or threatened in Illinois, because it was believed to have been extirpated (Brigham *et al.* 1981). This species, now known to exist in Illinois waters, has been proposed by the Illinois Endangered Species Technical Advisory Committee on Fishes for consideration as a threatened species. Additional studies, especially on the Mississippi and Kankakee rivers, are critical if the exact status of this fish in Illinois is to be determined.

## MUSSEL SURVEY

### Methods

Mussels were collected from the Kankakee River in Kankakee on 21 September 1984 for the purpose of determining the species composition and relative abundance of the mussel population at the site to be affected by the proposed FAU 6198 project (Figure 1). Collections were made under the supervision of INHS malacologist Jeanine M. Kasproicz, with the assistance of INHS biologists Mark J. Wetzell, Sherri L. Sandberg, Jeffrey T. Irish, Jean E. Karnes, Kevin S. Cummings, Patti L. Malmborg, and Glendy C. Vanderah. Approximately 30 man-hours of effort were expended during this mussel survey.

The study site was approximately 9,800 m<sup>2</sup> in area, extending to both banks of the river and to approximately 25 m upstream and 45 m downstream of a line bisecting the present Washington Avenue bridge. This area was sampled in sections roughly defined with respect to the bridge and each shoreline (Section 1 - south bank, downstream, near the Jobber's Park boatramp; Section 2 - mid-channel, downstream); Section 3 - north bank, downstream; Section 4 - south bank, upstream; Section 5 - mid-channel, upstream; and Section 6 - north bank, upstream (Figure 2). Each section was sampled by hand, in most cases as completely as possible. However, no attempt was made to collect all individuals from Section 5 (Figure 2), where mussels, primarily *Actinonaias carinata* and *Alasmidonta marginata*, were especially concentrated. The substrate in Section 6 was composed almost entirely of bedrock; no mussels

were collected from this area. Section H, an area surveyed for mussels in conjunction with a City of Kankakee project, will be discussed later in this report.

Mussels collected from each section were identified, enumerated, and returned live to the river. A few specimens were returned to the lab for positive identification. One or two individuals of most species were retained as voucher specimens. All retained specimens were deposited in the INHS mollusk collection.

### Results

During this survey 579 individuals representing 13 genera and 17 species of mussels were collected (Table 3), with *Actinonaias carinata* (Barnes) accounting for 70.3% of the collection. Other species which made up more than 1% of the collection included *Quadrula pustulosa* (Lea) (11.23%), *Alasmidonta marginata* Say (5.53%), *Megalonaias gigantea* (Barnes) (2.76%), *Cyclonaias tuberculata* (Rafinesque) (1.90%), *Lasmigona complanata* (Barnes) (1.21%), *Lasmigona costata* (Rafinesque) (1.21%), *Pleurobema cordatum* (Rafinesque) (1.21%), and *Ligumia recta* (Lamarck) (1.04%). In addition, two taxa of non-unionid bivalves also were documented: the Asiatic clam *Corbicula fluminea* (Muller) and the fingernail clam *Musculium* sp.

Riffles in and near the channel, with substrate composed of sand, gravel, rubble, and bedrock, and with a strong current, supported the greatest abundance of mussels, particularly *Actinonaias carinata*. However, the greatest diversity of mussel species was found in Section 1, where the substrate was generally finer and the current more gentle. Table 4 lists the number of individuals and species by section sampled (Figure 2).

### Discussion

The mussel community in the vicinity of the Washington Avenue bridge is one of the most abundant and diverse of any presently known from the Kankakee River. Seventeen species of unionid mussels were collected from the Washington Avenue bridge site for this survey on 21 September 1984. Including the results from another recent survey (Wetzel *et al.* 1984), 19 of the 20 unionid mussel species collected from the Kankakee River since 1976 were collected in September 1984 from within and immediately adjacent to this present study site.

Because sampling methods are difficult to standardize, results from various mussel surveys are not easily compared. Conditions such as depth, flow, water temperature substrate, and weather may be more or less favorable during different studies, and amount of effort expended per unit area sampled varies. Even when the effects of such variables are minimized, apparently random fluctuations in mussel populations at the same site have been observed from year to year (Van Cleve 1940). Nevertheless, some idea of the quality of the mussel community at the Washington Avenue bridge site can be obtained by comparing the results of other recent surveys, summarized below, with the findings of the present study. Extremely favorable sampling conditions were present during the Washington Avenue bridge survey on 21 September 1984.




Figure 2. Location of sections of the Kankakee River in the vicinity of the Washington Avenue bridge (U. S. routes 45 and 52), City of Kankakee, Kankakee County, Illinois, sampled for unionacean mussels on 21 September 1984.

Table 3. Species and numbers of mussels collected 21 September 1984 from the Kankakee River at the Washington Avenue bridge, City of Kankakee, Kankakee County, Illinois.

SPECIES	NUMBER
CLASS PELECYPODA	
Order Eulamellibranchia	
Superfamily Unionacea	
Family Unionidae	
<i>Actinonaias carinata</i> (Barnes)	407
<i>Actinonaias ellipsiformis</i> (Conrad)	5
<i>Alasmidonta marginata</i> Say	32
<i>Amblema plicata</i> (Say)	2
<i>Anodonta imbecillis</i> Say	5
<i>Cyclonaias tuberculata</i> (Rafinesque)	11
<i>Elliptio dilatata</i> (Rafinesque)	4
<i>Fusconaia flava</i> (Rafinesque)	1
<i>Lampsilis ovata</i> f. <i>ventricosa</i> (Barnes)	1
<i>Lasmigona complanata</i> (Barnes)	7
<i>Lasmigona costata</i> (Rafinesque)	7
<i>Ligumia recta</i> (Lamarck)	6
<i>Megalonaias gigantea</i> (Barnes)	16
<i>Pleurobema cordatum</i> (Rafinesque)	6
<i>Quadrula metanevra</i> (Rafinesque)	1
<i>Quadrula pustulosa</i> (Lea)	66
<i>Quadrula quadrula</i> (Rafinesque)	2
Superfamily Sphaeriacea	
Family Corbiculidae	
<i>Corbicula fluminea</i> (Muller)	+
Family Sphaeriidae	
<i>Musculium</i> sp.	+
Total	
579	

+ indicates presence of these taxa; actual numbers of individuals were not tabulated.

Table 4. Number of individuals and species of mussels collected 21 September 1984 from the Kankakee River at the Washington Avenue bridge, City of Kankakee, Kankakee County, Illinois.

Section	# of Individuals	# of species
1	147	16
2	193	9
3	122	8
4	86	10
5*	40+	5
6	-	-

\*Mussels, particularly *Actinonaias carinata* and *Alasmidonta carinata*, were especially concentrated in this section. No attempt was made to count all individuals of these two species collected from this section.

Five Kankakee River locations in Illinois were sampled in 1976 and 1978 by Lewis and Brice (1980). Eighteen species of live unionid mussels were collected in their study, 15 of which were present at their Custer Park site (27.2 km downstream of the Washington Avenue bridge). This site also was sampled quantitatively using SCUBA and wading (Brice and Lewis 1979), and yielded densities of 0 to 56 mussels/m<sup>2</sup>.

During her 1978 survey, Suloway (1981) sampled 13 river locations throughout the 90 km length of the Kankakee River in Illinois. Of these 13 stations, the site at the Washington Avenue bridge (Suloway's Station 6) yielded 12 of the 20 species collected during her entire survey. The number of individuals at this site (189) was second only to that taken from Suloway's Station 10 at Custer Park (197).

Thirteen species of unionid mussels were collected by INHS personnel during October 1979 from a stretch of the Kankakee River between Aroma Park (approximately 6.5 km upstream from the Washington Avenue bridge) and the Illinois-Indiana border (Brigham *et al.* 1981).

Brigham *et al.* (1984) recently conducted a mussel survey at a site 1.4 km downstream of the Washington Avenue bridge in connection with a City of Kankakee sewer interceptor project at Bird Park. Quantitative sampling of 36 0.5m<sup>2</sup> quadrats with SCUBA and wading revealed densities from 0 to 14 mussels/m<sup>2</sup>. Eleven species were collected.

Kasprowicz *et al.* (1985) surveyed the Kankakee River mussel fauna at Wilmington (28.8 km downstream of the Washington Avenue bridge) on 7 September 1984; 11 species were collected.

A survey of Kankakee River mussels was conducted on 27 September 1984 for the City of Kankakee (Wetzel *et al.* 1984) in conjunction with the reconstruction of a hydroelectric plant. This survey was conducted immediately downstream of the hydroelectric plant powerhouse draft tubes (Figure 2, Section H), an area adjacent to the Washington Avenue bridge project area. During the hydroelectric plant study, 33 individuals representing 6 genera and 7 species (*Actinonaias carinata*, *Anodonta grandis* Say, *Anodonta imbecillis*, *Lampsilis radiata siliquioidea* (Barnes), *Lasmigona complanata*, *Lasmigona costata*, and *Quadrula pustulosa*) were retrieved by a SCUBA diver from depths of 0.8 to 6.8 m. Two of these species (*Anodonta grandis* and *Lampsilis radiata siliquioidea*) were not collected during the Washington Avenue bridge study.

Nineteen species of unionacean mussels have been collected from the Kankakee River in the vicinity of the Washington Avenue bridge since 1978; 17 of these were collected during this present study. The mussel community at this site thus appears to be one of the most diverse, as well as one of the most abundant, of any known from the Kankakee River in Illinois.

## PRESENT STATUS OF MUSSELS IN THE KANKAKEE RIVER

Approximately 50 percent of the known species of freshwater mussels in the world occur in eastern North America (Stansbery 1970), achieving their greatest diversity in the Mississippi River basin (Johnson 1970). Significant depletions in both numbers of individuals and species have occurred in the last century throughout the country and have been documented for river systems such as the Illinois (Starrett 1971); Kankakee (Brigham *et al.* 1981; Suloway 1981); Kaskaskia (Suloway *et al.* 1981); Little Wabash (L. Suloway, unpublished); Ohio and Tennessee (Stansbery 1971); Rock (Miller 1972); Vermilion (Matteson and Dexter 1966; Suloway 1975); and the Wabash (Krumholz *et al.* 1970). A special case can be made for the protection of this unique fauna (Brigham *et al.* 1981).

In response to the decline in mussel populations, federal and state governments have enacted legislation to protect several species. The federal government currently lists 22 species of mussels found in the United States as endangered (U. S. Department of the Interior [USDI] 1984a). Lists of endangered threatened, rare, or protected mussels also have been established by several states adjacent to or near Illinois, including Alabama (Stansbery 1976); Arkansas (Sam Barkley, Arkansas Game and Fish Commission, pers. comm.), Indiana (Indiana Revised Statutes 1984), Kentucky (Babcock 1977, Branson *et al.* 1981); Michigan (van der Schalie 1975), Minnesota (Anon. 1984), Missouri (Wilson 1984); and Ohio and Wisconsin (McGaugh and Genoways 1976).

Illinois (Illinois Administrative Code 1984) recently emended their list of endangered and threatened species of vertebrates and plants to include nine federally endangered mussels presently occurring or known to formerly have occurred in Illinois waters. A number of freshwater mussels presently occurring or known to have occurred formerly in Illinois were proposed to the Endangered Species Technical Advisory Committee for state listing by INHS personnel on 26 March 1985 (Liane Suloway, pers. comm.), including *Plethobasus cyphus*, a species (one specimen) collected on 7 September 1984 from the Kankakee River at Wilmington in Will County (Kasproicz *et al.* 1985).

Studies during the last 100 years have yielded 37 species of mussels from the Kankakee River and its tributaries. Of these, only one, *Lampsilis higginsi* (Lea) (= *Lampsilis orbiculata* of Burch 1975), currently is considered federally endangered. Two other species reported from the Kankakee River, *Cumberlandia monodonta* (Say) and *Simpsoniconcha* (= *Simpsonaias*) *ambigua* (Say), currently are assigned to Category 2 (USDI 1984b), which means they are proposed for federal listing as endangered or threatened, but that conclusive data on biological vulnerability and threat currently are not available. These species have not been reported from the Kankakee River since 1906 (Suloway 1981). Eighteen of the 37 mussel species known to have occurred in the Kankakee River basin are listed as endangered, threatened, or rare in various states (Indiana, Kentucky, Michigan, Missouri, or Ohio). None of the 19 mussel species presently known to occur in the vicinity of the Washington Avenue bridge currently are listed as federally endangered or proposed for state listing.

The general decline in the mussel fauna of the Kankakee River in Illinois may be due to several factors, including pollution (domestic and industrial effluents and fertilizer and pesticide runoff), habitat destruction (dam construction, and increased sedimentation resulting from dredging and channelization, particularly in Indiana), and overharvest (commercial clamming, first during the early and mid-1900's for the pearl button industry, and recently for the cultured pearl industry in Japan). Only 20 species of freshwater mussels have been collected from the Kankakee River since 1978. Based upon available historic and current information, several species of mussels reported from the Kankakee River in Illinois presently could be considered uncommon or localized in Illinois and in some cases the United States. Two species found during this study, *Cyclonaias tuberculata* (Rafinesque) and *Ligumia recta* (Lamarck), are considered uncommon and on the decline in Illinois.

*Cyclonaias tuberculata* was collected by Matteson from the Fox, Kankakee, and the Vermilion (Wabash drainage) rivers during the 1950's. In Illinois, it recently has been collected from the Kankakee, Vermilion, Rock, and Ohio rivers but apparently is absent from the Des Plaines and Spoon rivers. It is uncommon and possibly extirpated from the upper Mississippi River.

*Ligumia recta* was collected by Matteson from the Fox and Kankakee rivers and the Rock River basin during the 1950's. This species was considered rare even at that time. This species recently has been collected from the Kankakee and Rock rivers, and also has been observed in the Mississippi (where it is rare) and the Ohio rivers in Illinois. Although this species once was common in the Illinois River, it apparently is extirpated now. This species also appears to be extirpated from the Kaskaskia and Spoon rivers in Illinois.

#### ACKNOWLEDGEMENTS

The authors wish to thank Survey biologists Patti L. Malmborg, Jeffrey T. Irish, Jean E. Karnes, Sherri L. Sandberg, and Glendy C. Vanderah for their field expertise. This project could not have been completed in a smooth and efficient manner without their assistance. Ms. Aleta Holt and Lloyd LeMere, INHS graphic artists, provided assistance with the maps. Dr. Lawrence M. Page provided valuable critique and Dr. Wallace E. LaBerge kindly reviewed this manuscript.

#### LITERATURE CITED

- Anon. 1984. Minnesota official list of endangered, threatened, and special concern plants and animals. January. [unpaged].
- Babcock, J. V. 1977. Endangered plants and animals of Kentucky. Univ. Kentucky Coll. Eng. Off. Res. Eng. Serv. Publ. 128 pp.
- Baker, F. C. 1906. A catalogue of the mollusca of Illinois. Bull. Illinois State Lab. Nat. Hist. 7:53-136.

- Becker, G. C. 1983. The fishes of Wisconsin. Univ. Wisconsin Press, Madison. xii + 1052 pp
- Branson, B. A., D. F. Harker, Jr., J. M. Baskin, M. E. Medley, D. L. Batch, M. L. Warren, Jr., W. H. Davis, W. C. Houtcooper, B. Monroe, Jr., L. R. Phillippe, and P. Cupp. 1981. Endangered, threatened, and rare animals and plants of Kentucky. Trans. Kentucky Acad. Sci. 42(3-4):77-89.
- Brice, J. R. and R. B. Lewis. 1979. Mapping of mussel (Unionidae) communities in large streams. Amer. Midl. Nat. 101(2):454-455.
- Brigham, A. R., L. B. Suloway, and L. M. Page. 1981. The effects of sedimentation on aquatic life of the Kankakee River. Phase II: Quantitative studies and threatened, endangered, and rare species. Illinois Dept. Energy Nat. Resour. Document No. 81/37. ii + 16 pp.
- Brigham, W. U., L. Suloway, J. M. Kasprowicz, and M. J. Wetzel. 1984. Survey of Kankakee River mussels (Mollusca: Unionidae) at site of proposed sewer interceptor project, City of Kankakee, Kankakee County, Illinois. [A report prepared for the City of Kankakee Department of Water Pollution Control, dated 27 August] ii + 12 pp.
- Burch, J. B. 1975. Freshwater unionacean clams (Mollusca: Pelecypoda) of North America. Revised edition. Malacological Publications, Hamburg, MI. xviii + 204 pp.
- Cahn, A. R. 1927. An ecological study of southern Wisconsin fishes. The brook silversides (*Labidesthes sicculus*) and the cisco (*Leucichthys artedii*) in their relations to the region. Illinois Biol. Monogr. 11(1):1-151.
- Clay, W. M. 1975. The fishes of Kentucky. Kentucky Dept. of Fish and Wildlife Resources, Frankfort. 416 pp.
- Clemmer, G. H. 1980. *Notropis amnis* (Hubbs and Greene), pallid shiner. p. 224, In D. S. Lee *et al.*, Atlas of North American freshwater fishes. North Carolina State Museum of Natural History, Raleigh. 854 pp.
- Cross, F. B. 1967. Handbook of fishes of Kansas. Univ. Kansas Mus. Nat. Hist. Misc. Publ. 45. 357 pp.
- Eddy, S., and J. C. Underhill. 1974. Northern fishes with special reference to the Upper Mississippi Valley. 3rd edition. University of Minnesota Press, Minneapolis. 414 pp.
- Forbes, S. A., and R. E. Richardson. 1920. The fishes of Illinois. 2nd ed. Illinois St. Lab. Nat. Hist., Urbana. cxxxvi + 357 pp.
- Gilbert, C. H. 1980. *Notropis heterolepis* Eigenmann and Eigenmann, blacknose shiner. p. 272, In D. S. Lee *et al.* Atlas of North American freshwater fishes. North Carolina St. Mus. Nat. Hist., Raleigh.

- Harlan, J. R., and E. B. Speaker. 1969. Iowa fish and fishing. State Conservation Commission, Des Moines. 365 pp.
- Hubbs, C. L. 1951. *Notropis amnis*, a new cyprinid fish of the Mississippi fauna, with two subspecies. Occasional Papers of the Museum of Zoology, University of Michigan 530:1-31.
- Illinois Administrative Code. 1984. Title 17: Conservation. Chapter I: Department of Conservation. Subchapter c: Endangered species. Part 1010. Illinois list of endangered and threatened fauna. Sections 1010.10 through 1010.40. July 25, 1984.
- Indiana Revised Statutes. 1984. Discretionary order no. W-12, revised August 1984. Endangered, threatened, and special concern wildlife. Section 6. 310 IAC 3-3-6, amended. [pers. comm. from James M. Ridenour, Director, Indiana Department of Natural Resources] [5 pp.]
- Ivens, J. L., N. G. Bhowmik, A. R. Brigham, and D. L. Gross. 1981. The Kankakee River yesterday and today. Illinois State Water Survey, Miscellaneous Publication 60. 24 pp.
- Johnson, R. I. 1970. Systematics and zoogeography of Unionidae of the southern Atlantic slope region. Bull. Mus. Comp. Zool. 140:263-449.
- Kasprowicz, J. M., M. J. Wetzel, K. S. Cummings, and W. U. Brigham. 1985. Survey of Kankakee River fishes and mussels at Wilmington/Illinois Route 53 bridge, City of Wilmington, Will County, Illinois. IDOT Job Number P-91-051-79. [A draft report prepared for the Illinois Department of Transportation Bureau of Location and Environment, 2300 South Dirksen Parkway, Springfield, Il 62764. Date: 6 February. ii + 26 pp.]
- Krumholz, L. A., R. L. Bingham, and E. R. Meyer. 1970. A survey of the commercially valuable mussels of the Wabash and White Rivers of Indiana. Proc. Indiana Acad. Sci. 79:205-226.
- Lewis R. B., and J. R. Brice. 1980. A comparison of the past and present freshwater mussel fauna of the Kankakee River in Illinois. Chicago Acad. Sci. Nat. Hist. Misc. 211. 7 pp.
- Lopinot, A. L., and P. W. Smith. 1973. Rare and endangered fish of Illinois. Illinois Dept. Conserv., Springfield. 53 pp.
- Matteson, M. R., and R. W. Dexter. 1966. Changes in pelecypod populations in Salt Fork of Big Vermilion River, Illinois, 1918-1962. Nautilus 79:96-101.
- McGaugh, M. H., and H. H. Genoways. 1976. State laws as they pertain to scientific collecting permits. Texas Tech. Univ. Mus. Sci. Program. Museology No. 2. 81 pp.
- Miller, R. J., and H. W. Robison. 1973. The fishes of Oklahoma. Oklahoma State University Press, Stillwater. 246 pp.

- Miller, T. B. 1972. Investigation of the freshwater mussels of the Rock River, Illinois. Illinois Dept. Conserv., Div. Fish Spec. Rept. No. 43. 12 pp.
- Nurnberger, P. K. 1928. A list of the plant and animal food of some fishes of Jay Cooke Park. Trans. Amer. Fish. Soc. 58:175-177.
- Nordstrom, G. R., W. L. Pflieger, K. C. Sadler, and W. H. Lewis. 1977. Rare and endangered species of Missouri. Missouri Dept. Conserv. and U. S. Dept. Agric. Soil Conserv. Serv. 129 pp.
- Page, L. M., P. W. Smith, L. B. Suloway, A. R. Brigham, D. W. Webb, J. D. Unzicker, and W. U. Brigham. [1979]. The effects of sedimentation on aquatic life of the Kankakee River. Phase I. Illinois Nat. Hist. Surv. 86 pp.
- Parmalee, P. W. 1967. The fresh-water mussels of Illinois. Illinois St. Mus. Pop. Sci. Ser. Vol. 8. 108 pp.
- Pflieger, W. L. 1975. The fishes of Missouri. Missouri Dept. Conserv., Box 180, Jefferson City, MO 65101. viii + 343 pp.
- Scott, W. B., and E. J. Crossman. 1973. Freshwater fishes of Canada. Fish. Res. Bd. Canada Bull. 184. Ottawa. xviii + 966 pp.
- Skelly, T. M., and M. J. Sule. 1980. Water quality of the Kankakee River in the Braidwood aquatic monitoring area, 1979. Section 8, In Construction and Preoperational Aquatic Monitoring Program for the Kankakee River. Braidwood Station Third Annual Report. Illinois Natural History Survey, Champaign, Illinois.
- Skelly, T. M., and M. J. Sule. 1983. The pallid shiner, *Notropis amnis* Hubbs and Greene, a rare Illinois fish. Trans. Illinois St. Acad. Sci. 76(1&2):131-140.
- Smith, P. W. 1979. The fishes of Illinois. Univ. Illinois Press Urbana. xxix + 314 pp.
- Smith, P. W., and L. M. Page. 1981. Endangered and threatened fishes. pp. 5-20, In Natural Land Institute. Endangered and threatened vertebrate animals and vascular plants of Illinois. Illinois Dept. Conserv., Springfield.
- Stansbery, D. H. 1970. Eastern freshwater mollusks, the Mississippi and St. Lawrence River systems. pp. 9-21, In A. H. Clarke, ed. Papers on rare and endangered mollusks of North America. Malacologia 10:1-56.
- Stansbery, D. H. 1976. Naiad mollusks. pp. 42-52, In H. Boschung, ed. Endangered and threatened plants and animals of Alabama. Bull. Alabama Mus. Nat. Hist. No. 2. Univ. of Alabama, University, AL.
- Starrett W. C. 1971. A survey of the mussels (Unionacea) of the Illinois River: a polluted stream. Illinois Nat. Hist. Surv. Bull. 30(5):267-403.

- Suloway, J. J. 1975. Changes in the molluscan populations of the Salt Fork of the Big Vermilion River since F. C. Baker's study of 1918-1920. Unpubl. M.S. thesis, Dept. Biol., Univ. of Illinois, Urbana. 52 pp.
- Suloway, L. 1981. The unionid (Mollusca: Bivalvia) fauna of the Kankakee River in Illinois. *Amer. Midl. Nat.* 105(2):233-239.
- Suloway, L., J. J. Suloway, and E. E. Herricks. 1981. Changes in the freshwater mussel (Mollusca: Pelecypoda: Unionidae) fauna of the Kaskaskia River, Illinois with emphasis on the effects of impoundment. *Trans. Illinois St. Acad. Sci.* 74(1+2):79-90.
- Sweet M. 1984. Office memorandum to members, ESTAC [Endangered Species Technical Advisory Committee] on fish. Illinois Department of Conservation. 6 March. unpagged.
- Trautman, M. B. 1957. The fishes of Ohio. Ohio St. Univ. Press, Columbus. xvii + 683 pp.
- U. S. Department of the Interior, Fish and Wildlife Service (USDI). 1984a. Box score of listings/recovery plans. *Endangered Species Tech. Bull.* 9(6):12.
- U. S. Department of the Interior, Fish and Wildlife Service (USDI). 1984b. 50 CFR Part 17. Endangered and threatened wildlife and plants; Review of invertebrate wildlife for listing as endangered and threatened species. *Federal Register* 49(100)(Part III):21664-21675.
- Van Cleve, H. J. 1940. Ten years of observation on a fresh-water mussel population. *Ecology* 21(3):363-370.
- Van der Schalie, H. 1975. An ecological approach to rare and endangered species in the Great Lakes region. *Michigan Acad.* 8:7-22.
- Wetzel, M. J., J. M. Kasprowicz, and W. U. Brigham. 1984. Survey of Kankakee River mussels (Mollusca) at site of proposed hydroelectric plant reconstruction, City of Kankakee, Kankakee County, Illinois. [A report currently in preparation for the City of Kankakee Department of Water Pollution Control].
- Wilson, C. B., and H. W. Clark. 1912. The mussel fauna of the Kankakee basin. U.S. Dept. Commerce and Labor, Bureau of Fisheries, Doc. No. 758. 52 pp.
- Wilson, J. H., ed. 1984 [1985]. Rare & endangered species of Missouri. Missouri Dept. Conserv. P.O. Box 180, Jefferson City, MO 65102. 171 pp.