

BROADBAND MICROWAVE AND COMPUTATIONAL STUDY OF HEXAFLUORO-O-XYLENE: HIGHLY COUPLED CF₃ ROTORS

SVEN HERBERS, SEAN FRITZ, PIYUSH MISHRA, YONGBIN KIM, LYUDMILA V SLIPCHENKO, TIMOTHY S. ZWIER, *Department of Chemistry, Purdue University, West Lafayette, IN, USA.*

The rotational constants and quartic centrifugal distortion coefficients of hexafluoro-o-xylene and all singly ¹³C isotopologues were precisely determined from the 8 to 18GHz gas phase microwave spectrum. A preliminary r₀ structure was determined, reproducing the experimental rotational constants with deviations of no more than 15kHz. Interestingly, rather than the C_{2v} symmetry structure expected intuitively, as in o-xylene, calculations with a variety of methods (B3LYP, CAM-B3LYP, ωB97XD, MP2, and CCSD(T)) predict a C₂ symmetry structure in which the two CF₃ groups rotate in opposite directions by about 16 degrees. Analysis of the interactions between the two CF₃ groups using an effective fragment potential (EFP) approach identified two major contributions to their interaction, due to exchange repulsion and electrostatic repulsion, with electrostatic repulsion responsible for the barrier at the C_{2v} geometry.

SH, SMF, PM and TSZ gratefully acknowledge support for this work from the Department of Energy Basic Energy Sciences Gas Phase Chemical Physics program under Grant No. DE-FG02-96ER14656. YK and LVS gratefully acknowledge support from the National Science Foundation (NSF CHE-1800505). Present address for TSZ: Combustion Research Facility, Sandia National Laboratory, Livermore, CA 94550.