



LIBRARY
OF THE
UNIVERSITY
OF ILLINOIS

cop.2

NATURAL HISTORY
SURVEY

64
STATE OF ILLINOIS
DEPARTMENT OF REGISTRATION AND EDUCATION
NATURAL HISTORY SURVEY DIVISION

The Value of IN VITRO FUNGICIDE TESTS

DAN NEELY

Illinois Natural History Survey
Biological Notes No. 64

Urbana, Illinois January, 1969

The Value of IN VITRO FUNGICIDE TESTS

Dan Neely

Because chemists now seem able to synthesize an unlimited number of compounds, we can expect that many fungicides developed in the future will prove to be better than those we are using at the present time. Appropriate methods must be employed to select the better fungicides from among the mass of candidate materials.

Separating the successful and unsuccessful fungicides is an expensive procedure. Wellman (1967) estimated industry's cost of research on one new fungicide to be \$3 million. Since it becomes increasingly difficult to find a fungicide that is more successful than one just introduced, these costs can be expected to increase. With this vast expense involved, research agencies must develop and use techniques and methods that are both efficient and reliable.

The three traditional steps in the search for an effective new fungicide are laboratory evaluation, greenhouse evaluation, and field evaluation. With the proper tests and the proper interpretation of the test results, this still seems to be the only practical system. Opinions vary among investigators as to the importance of each of the steps. This paper will review the test methods used to seek and evaluate fungicides in the laboratory, and specifically the *in vitro* tests. It will be further limited to those methods used to develop protective foliar fungicides. The specialized methods used to develop eradicated, systemic, and soil fungicides are not included.

The value of *in vitro* tests to appraise fungicides in the laboratory is determined by the ease, speed, and simplicity of test manipulations and the reliability and usability of test results. Alternative *in vivo* methods are time consuming, which prohibits extensive use of greenhouse and field techniques in the initial evaluation of chemicals. *In vivo* tests are essential, however, in determining how effectively chemicals control plant diseases.

The literature concerning *in vitro* testing of fungicides has been reviewed by several investigators (Horsfall 1945a, 1945b, 1956; McCallan 1947, 1959), most recently by Torgeson (1967). In this paper I wish to emphasize the variety of *in vitro* techniques that can be used as the basis for selecting fungicides

for greenhouse and field studies. Many of these *in vitro* techniques have additional value, for in recent years they have been extensively used in studies seeking the toxic principles of fungicide groups and the mode of action of chemical structures on plant pathogens.

The significance of fungicide tests in an artificial rather than a natural situation is occasionally questioned. Marsh (1936) and Howard (1939) compared spore germination on leaves and on glass slides and found comparable results. Miller (1943) compared the retention of copper on leaves and on pyralin plates and found laboratory methods to be fairly accurate.

FUNGICIDE-FUNGUS INTERACTIONS

In vitro tests are used to measure and rank the fungitoxicity of fungicides, to measure fungal specificity, and to measure fungicide deposition, redistribution, tenacity, persistence, stability, and volatility.

Chemicals toxic to fungi have been observed to affect the fungi in various ways. The fungus mycelium may cease growing, change metabolic processes, or be killed. The fungus spores may fail to swell before germinating, fail to germinate, plasmolyze, or be killed. Each of these reactions is measurable. Each has been used in the past to determine fungicide toxicity *in vitro*. Standard methods of determining toxicity have not been universally accepted.

Fungitoxicity

While we accept the general definition of a fungicide as "any substance that destroys fungi or inhibits the growth of the spores or hyphae," we should restrict the use of the adjective *fungicidal* to the conditions that kill fungi and the term *fungistatic* to those conditions that prevent additional growth or sporulation of fungi without killing them (McCallan & Wellman 1942). The term *fungitoxic* has come into use in those situations where detrimental morphological or physiological fungal changes occur but no effort is made to determine whether or not the fungi are killed by the treatment.

Many techniques have been developed during the last 40 years that attempt to measure the fungitoxicity of chemicals or compare the fungitoxic values of chemicals. These techniques are usually bioassays involving a chemical and a test fungus. The use of a fungus introduces biological variability and reduces

This paper is published by authority of the State of Illinois, IRS Ch. 127, Par. 58.21, and is a contribution from the Section of Botany and Plant Pathology of the Natural History Survey. It was earlier presented in England, in a symposium on methods of testing fungicides, at the First International Congress of Plant Pathology, London, July 14-26, 1968. Dr. Dan Neely is a plant pathologist at the Survey.

the significance of each measurement of toxicity. Many of the techniques now used have been developed with the express purpose of reducing biological variability.

Testing With Spore Germination.—Early tests of Reddick & Wallace (1910) and McCallan and co-workers (McCallan 1930; McCallan, *et al.* 1941*b*; McCallan & Wilcoxon 1938, 1939; Wilcoxon & McCallan 1939) and Horsfall, *et al.* (1940) were spore germination tests involving the deposition of chemicals on glass slides, evaporation to dryness, and the addition of a drop of water containing spores of the test fungus. The American Phytopathological Society (1943) published instructions for this technique.

One problem encountered with this technique was the variation in size of the water drops. Montgomery & Moore (1938) attempted to solve this by drawing a circle on the glass slide with a diamond pencil, Young (1944) etched a circle with hydrofluoric acid, and Miller (1949) drew a circle with a wax pencil. Others have used depression slides (Barratt & Horsfall 1947, Kirby & Frick 1953, and Prusova 1962), small petri dishes (Tamura 1954), microbeakers (Shafer 1952 and Spencer 1962), or raised cover slips on slides (Peterson 1941).

Another problem was that of obtaining an accurate determination of the amount of toxic material on the slide. Elaborate laboratory sprayers have been built to uniformly deposit the chemical. An alternate solution by McCallan & Wilcoxon (1938) and Peterson (1941) involved mixing the toxicant with the spore suspension directly and not previously forming a dry deposit on the slide. Procedures for this test tube dilution technique were described by the American Phytopathological Society (1947).

Although germinating spores on glass slides are easily examined microscopically, glass is not the ideal substrate for spore germination. Marsh (1936) observed that fungus spores germinated better on glass slides that had been coated with cellulose. Transparent materials other than glass have been successfully used. McIntosh (1961) and Spencer (1962) used transparent plastic polymers. Neely & Himelick (1966) germinated spores on nonwaterproof cellophane discs placed on filter paper pads previously saturated with the toxic solution or suspension. The cellophane discs were transferred to slides for microscopic determination of fungistatic activity or transferred to agar for determination of fungicidal activity.

Many fungus spores do not germinate well in water drops. Davies, *et al.* (1948) found that shaking the culture greatly improved the uniformity of spore germination. Darby (1960) described a shaker method with a 4-hour germination period. Cattani (1954) observed that spores germinate better on agar than in water drops. He mixed the fungicide with warm agar, allowed the agar to solidify, seeded the

agar with spores, and counted the germinating spores by placing the petri dish under a microscope. Heyns, *et al.* (1965) placed seeded, toxicant-agar preparations on microscope slides.

Rapid tests involving swelling of spores prior to spore germination also have been used to measure fungitoxicity. Mandels & Darby (1953) described a 3-hour test with the failure of spores to swell as the measurable character. Koopmans (1959) based his results on the loss of turgidity of powdery mildew spores in toxic solutions.

Macroscopic Tests.—Counting germinating spores has always been a tedious, fatiguing task. In vitro tests utilizing macroscopic observations of fungus colony growth have often been used. Lee & Martin (1927) and Wellman & Heald (1940) placed fungus spores in known aqueous concentrations of the toxicant for varying lengths of time before transferring them to bouillon or agar. Their tests are based on reduction in number of fungus colonies formed.

The most commonly used macroscopic tests involve germination of spores on toxicant-agar preparations in petri plates with colony counts being made or the absence of colonies noted (Palmiter & Keitt 1937). With the roll culture technique of Manten, *et al.* (1950), the toxicant-agar layer is formed along the sides of round bottles rather than in petri dishes. A closely related method involves placing known quantities of toxicant at specific spots on seeded agar plates. The fungicide diffuses through the agar. The diameter of the clear area where spore germination was inhibited is measured. The fungicide may be placed in holes in the agar (Mildner, *et al.* 1963) or added with saturated filter paper pads (Thornberry 1950 and Leben & Keitt 1950) or string (Kuhfuss 1957).

In vitro techniques measuring toxicity of chemicals to fungus hyphae are often useful, especially with fungi that fail to sporulate or sporulate poorly in the laboratory. A common procedure is to incorporate toxic materials into agar and then seed with uniform mycelial colonies in agar blocks (Carpenter 1942), on filter paper pads (Sharville & Pelletier 1956), or on agar-coated cover slips (Bomar 1962). The reduction in fungus growth is then measured. Fungus cultures on agar plates also have been dusted or sprayed with toxic materials (Henry & Wagner 1940), submerged beneath toxic materials (Grosser & Friedrich 1947), or placed near toxic materials (Moreau & Moreau 1959), after which the resulting inhibition of growth was measured.

The fungus mycelium growing in toxic solutions has been linearly measured (Mason & Powell 1947) and weighed (Le Tourneau & Buer 1961). Forsberg (1949) placed a fungus-infested string in toxic dilutions for varying periods of time, then transferred the string to agar to determine fungicidal activity.

Mandels & Sin (1950) describe fungitoxicity tests

with a manometric system where the growth of the organism is measured by the amount of oxygen absorbed. McCallan, *et al.* (1954) and McCallan & Miller (1957) felt that measurement of oxygen consumption was not a particularly useful means of determining fungitoxicity.

Comparisons of Methods of Testing.—There have been relatively few published reports in which the methods for determining fungitoxicity have been compared. In observations of techniques involving spore germination, Frick (1964) found the test tube dilution test less variable than the dried deposit-slide germination test. Gottlieb (1945) found the seeded, toxicant-agar method and the test tube dilution method equally sensitive. Manten, *et al.* (1950) found the roll culture method more useful than the seeded toxicant-agar, dried deposit-slide germination, and gravimetric methods. Himelick & Neely (1965) found the cellophane transfer technique more sensitive than the seeded agar-toxicant spot method. Walker (1955) compared spore germination, spore respiration, and mycelia respiration techniques and found that, in general, the three methods gave the same ranking of fungicides.

Measurements.—In establishing the value of fungitoxicity tests we must not only know the method of testing but also the method of measurement and the means of expressing the resulting data. In bioassays this is a difficult task. McCallan and his co-workers have found that the amounts of fungicides absorbed by spores of different fungal species vary greatly and suggest that possibly fungitoxicity should not be based on the concentration of the external solution or suspension but on the weight of toxicant actually absorbed by the fungus (McCallan & Miller 1958; Miller, *et al.* 1953).

One problem encountered with the dried deposit-slide germination technique was whether to express the results in terms of weight of toxicant per unit area of slide or weight of toxicant per unit volume of water. Since the amount of toxicant in solution was not readily known, the results were usually expressed in weight per unit area.

A second problem in giving the results with spore germination tests was how to express the percentage of germination at different fungicide concentrations. It could be shown graphically with ease using a dosage-response curve, but it was difficult to express verbally. McCallan & Wilcoxon (1938) used the term LD₅₀ for that toxic concentration at which there was a reduction of 50 percent in spore germination. Later they used logarithmic paper to estimate the LD₅₀ (Wilcoxon & McCallan 1939), more appropriately called the ED₅₀ (McCallan 1948).

Interpretations of the role of the dosage-response curve in the evaluation of fungicides and the increasingly elaborate statistical methods that accompanied

use of the dosage-response curve were the subjects for numerous research papers from 1940 to 1960 (Dimond, *et al.* 1941; Horsfall 1956; Litchfield & Wilcoxon 1949; McCallan, *et al.* 1959). In many reports written during this period the results were expressed in a statistical language unreadable by many plant pathologists. Kundert (1956) commented on the exaggerated application of statistics in the assay of fungicides. Many researchers are now reporting results as the lowest external concentration that completely inhibits spore germination or mycelial growth expressed in parts per million or its equivalent (Guillemat & Lambert 1960; Pianka, *et al.* 1966; Luijten & van der Kerk 1961; Neely & Himelick 1966; Nisikado, *et al.* 1951; Pluijgers & Kaars Sijpesteijn 1966).

Fungal Selectivity

Although phycomycetes, powdery mildew fungi, other ascomycetes, and basidiomycetes are not equally sensitive to fungicides and although certain fungi are sensitive or resistant to specific fungicides (Horsfall 1951; Horsfall & Lukens 1966; Wellman & McCallan 1943), most fungicides now in commercial use have a broad spectrum of activity against fungal species. The test organism commonly used in the laboratory will often rank candidate fungicides in the same order as a specific disease-causing organism. McCallan, *et al.* (1941a), in a spore germination test with 6 fungi and 20 compounds, reported the fungi similarly sensitive. Neely & Himelick (1966), in a test with 7 fungi and 24 compounds, in general confirmed these results. Casarini & Pucci (1957) in a test with 6 fungi and 6 compounds felt the resulting differences between fungi sufficiently variable to recommend that the same fungus be used in *in vitro* trials as in field trials whenever possible.

Deposition, Redistribution, Tenacity, Stability

To be successful a fungicide must prove itself on a natural surface and in a natural environment. This involves the physical and chemical factors of deposition, redistribution, tenacity, and stability (Burchfield 1960, 1967). Modifications of the basic fungitoxicity methods have been developed with the aim of learning as much as possible about these additional factors in the laboratory.

Redistribution of a fungicide is often essential for control of a disease because plant growth, poor tenacity, or poor coverage may leave the fungicide particles widely distributed. Powell (1961) measures the ability of fungicides to redistribute by placing a known fungicide concentration in one depression of a glass slide and a spore suspension in the second de-

pression, making a water bridge between the two with filter paper, and measuring the decrease in spore germination.

The tenacity of fungicide deposits on glass slides was measured by Heuberger (1940) by dipping and swishing the slides several times in water, allowing them to dry, and seeding with a spore suspension. Chapman, *et al.* (1950) suspended the slides in circulating water and removed them at varying intervals of time before drying and seeding. Kovacs (1961) applied the fungicides on polyethylene discs, sprayed them with artificial rain, and placed the discs on seeded agar.

The stability of fungicide deposits on glass slides was measured by Barratt (1946) by subjecting them to high humidity and seeding with spore suspensions. Serra (1964) exposed polyethylene discs dusted with fungicides to ultraviolet light for varying periods of time, then transferred the discs to seeded agar. Ciferri, *et al.* (1961) found that zineb frequently increased in fungitoxicity after 2 to 6 years storage although chemical tests showed a decrease in the amount of "active ingredient."

Predicting Field Performance

In vitro fungitoxicity studies are often used as a means of predicting the field performance of candidate fungicides. The literature relating to predicting field performance from laboratory tests has been reviewed by Ciferri (1952), Rich, *et al.* (1953), McCallan (1959), Block (1959), and Torgeson (1967).

Horsfall and his co-workers were among the first to state that the protective value of fungicides could be predicted on the basis of laboratory results. Horsfall, *et al.* (1941) felt that fungitoxicity and tenacity were the two fundamental components of protective value. Keil, *et al.* (1952) called attention to the necessity of chemical stability. Waggoner, *et al.* (1952) later stated that a statistical prediction of field performance could be made on the basis of toxicity, tenacity, and stability of fungicides.

The relationships between laboratory and field results are not always clear cut. A hesitancy to predict field results from laboratory tests often arises in conversations or personal correspondence, but is infrequently found in the literature. Martin (1942), however, stated that to expect a simple correlation between fungitoxicity alone and field performance is too ambitious. Smith & Read (1961), in studies with the cucumber powdery mildew fungus, found a great variance between laboratory and field results.

Yet, in vitro bioassays often correlate well with field trials. Holloman & Young (1951) reported that laboratory assay was a reliable criterion for field performance for control of *Botrytis* leaf spot on

gladioli. Klomprens & Vaughn (1952) reported field and laboratory trials very consistent for control of *Helminthosporium* on bent grass. Brook (1957) found that laboratory and greenhouse experiments rated fungicides in almost the same order for control of *Botrytis* gray mold on tomatoes. Monroe (1963), in studies on bean powdery mildew and bean rust, stated that in vitro and in vivo methods generally ranked the fungicides in the same order. Misato (1963), in studies on rice blast, found the in vitro methods measuring inhibition of mycelial growth and spore formation to be satisfactorily correlated with field results.

Although a prediction based on in vitro tests may prove erroneous on a particular host with a particular set of environmental conditions in the field, the success or failure of the fungicide can often be explained with confidence if a sufficiently varied group of in vitro tests have been performed. Most researchers seeking protective fungicides at least agree with the statement that in vitro testing will result in the selection of chemicals worthy of greenhouse and field testing.

FUNGICIDE-FUNGUS-HOST INTERACTIONS

In vitro tests in the laboratory will not in the foreseeable future replace greenhouse and field trials (Hamilton 1959 and McCallan 1966). An effort is being made, however, to introduce a plant or plant part into in vitro tests so that the reactions of fungicide, fungus, and plant are present. Leben (1949) sprayed leaf discs with an antibiotic, then transferred the leaf discs to seeded agar to determine if the material remained fungitoxic. Leben & Keitt (1949) applied artificial rain to fungicide-treated leaf discs to see if the material was tenacious. Chaves (1954) evaluated the effects of fungicide stickers in much the same manner. Adams, *et al.* (1951) devised a leaf punch so that numerous leaf samples with uniform areas could be easily obtained for use in fungicide bioassays.

The ultimate step in in vitro testing is not simply a bioassay using a plant part, but the prevention of the disease itself. Schmidt (1951) treated the primary leaves of celery with fungicides, inoculated with *Septoria apii*, and obtained results on protective fungicides within 2 weeks. The laboratory fungicide trials of Hislop & Park (1962), using detached, treated, and inoculated pods of *Theobroma cacao*, correlated well with field tests for control of *Phytophthora palmivora*. Horn (1964) described a similar detached cucumber leaf method for screening fungicides for control of cucumber anthracnose. Niemann & Dekker (1966) described a method of evaluating compounds for control of powdery mildew of cucumber by floating leaf discs treated with fungicides on water, inoculating, and observing disease symptoms.

CONCLUSIONS

In vitro fungicide tests are extensively used in the initial evaluation of potentially valuable fungicides. In recent years these tests have provided valuable information on the toxicity of various chemical structures and their modes of action against fungi.

The methods of testing to determine the fungitoxicity of chemicals are extremely varied and are constantly changing. This probably means that an acceptable method that can be universally used as a standard procedure is yet to be developed. The procedures for fungicide testing should be quick, easy, and simple and the results should be reliable, reproducible, and usable.

In vitro fungicide tests are used not only to determine fungitoxicity of test chemicals but also range and degree of susceptibility of fungi, deposition and redistribution of fungicides, stability to resist chemical

change, tenacity to resist rainfall and wind, and persistence to resist other degrading elements. The results from these in vitro tests often will explain why a fungicide performed well or poorly in the field.

When a varied group of in vitro tests are performed in the laboratory, the candidate fungicides most likely to give satisfactory protection to plants will be selected for continued greenhouse and field testing. Not all of the complex chemical and environmental conditions of disease development can be reproduced either in the laboratory or the greenhouse. The introduction of plants or plant parts into in vitro tests does more closely approach natural conditions, and the use of these techniques will probably increase in the future.

At present there are no practical substitutes for in vitro tests for the initial screening of candidate fungicides. Thus these tests are invaluable as one of a series of steps required in the evaluation of fungicides.

LITERATURE CITED

- ADAMS, ROBERT E., ROBERT M. PRATT, and C. W. TERRY. 1951. A leaf punch for sampling foliage for analysis of spray deposits. *Phytopathology* 41:568-569.
- AMERICAN PHYTOPATHOLOGICAL SOCIETY, COMMITTEE ON STANDARDIZATION OF FUNGICIDAL TESTS. 1943. The slide-germination method of evaluating protectant fungicides. *Phytopathology* 33:627-632.
- . 1947. Test tube dilution technique for use with the slide-germination method of evaluating protectant fungicides. *Phytopathology* 37:354-356.
- BARRATT, R. W. 1946. A laboratory assay for stability of organic fungicide residues. *Phytopathology* 36:679. (Abstr.)
- , and JAMES G. HORSFALL. 1947. Fungicidal action of metallic alkyl bisdithiocarbamates. *Connecticut Agricultural Experiment Station Bulletin* 508. 51 p.
- BLOCK, S. S. 1959. Getting the most from fungicide tests. *Journal of Agricultural and Food Chemistry* 7:18-23.
- BOMAR, M. 1962. Estimation of efficiency of fungitoxic compounds according to the inhibition of mycelium growth. *Folia Microbiologica* 7:185-190.
- BROOK, P. J. 1957. A comparison of glasshouse and laboratory methods for testing fungicides against *Botrytis cinerea*. *New Zealand Journal of Science and Technology* 38(A):506-511.
- BURCHFIELD, H. P. 1960. Performance of fungicides on plants and in soil—Physical, chemical, and biological considerations, p. 477-520. In J. C. Horsfall and A. E. Dimond [ed.] *Plant Pathology*, Vol. III. Academic Press, Inc., New York.
- . 1967. Chemical and physical interactions, p. 463-508. In Dewayne C. Torgeson [ed.] *Fungicides*, an advanced treatise, Vol. I. Academic Press, Inc., New York.
- CARPENTER, J. B. 1942. A toximetric study of some eradicant fungicides. *Phytopathology* 32:845-856.
- CASARINI, BRUNO, and ENRICO PUCCI. 1957. Prove "in vitro" della efficacia fungicida di diversi antierittogamici su funghi scelti fra gli abituali "test" e fra alcuni dannosi parassiti delle piante. *La Ricerca Scientifica* 27:1468-1477.
- CHAPMAN, R. A., JAMES G. HORSFALL, and H. L. KEIL. 1950. Predicting protective value of fungicides from slope of wash-off curve. *Phytopathology* 40:4-5. (Abstr.)
- CHAVES, GERALDO M. 1954. Método bioanalítico para evaluación comparativa de adherentes en fungicidas orgánicos. *Turrialba* 4:39-42.
- CIFERRI, R. 1952. Revisione dei metodi di controllo biologico proposti per gli anti erittogamici e metodi adottati in Italia. *Istituto Botanico della Università Laboratorio Crittogamico, Pavia, Atti Serie 5*, 10:3-23.
- , A. SCARDAVI, and P. TRIVELLI RICCI. 1961. Modificazioni del potere antifungino dei formulati a base di zinco dopo due a sei anni d'immagazzinamento. *Notiziario sulle Malattie delle Piante* 56(NS35):59-72.
- DARBY, RICHARD T. 1960. Fungicide assay by spore germination in shaker flasks. *Applied Microbiology* 8:146-148.
- DAVIES, O. L., R. B. DUCKWORTH, and G. C. M. HARRIS. 1948. A method for estimating percentage germination of fungal spores. *Nature* 161:642.
- DIMOND, ALBERT E., JAMES G. HORSFALL, J. W. HEUBERGER, and E. M. STODDARD. 1941. Role of the dosage-response curve in the evaluation of fungicides. *Connecticut Agricultural Experiment Station Bulletin* 451:635-667.
- FORSBERG, J. L. 1949. A new method of evaluating fungicides. *Phytopathology* 39:172-174.
- FRICK, E. L. 1964. Methods of reducing variability in the results of glass-slide spore-germination assays of fungitoxicity. *Annals of Applied Biology* 54:349-360.
- GATTANI, M. L. 1954. The agar plate spore germination method for testing fungicides. *Phytopathology* 44:113-115.
- GOTTLIEB, DAVID. 1945. A comparison of the agar plate and test-tube-dilution methods for the preliminary evaluation of fungicides. *Phytopathology* 35:485. (Abstr.)
- GROSSER, ANNEMARIE, and WILHELM FRIEDRICH. 1947. Eine testmethode zur Bestimmung von Pilzhemmstoffen. *Zeitschrift für Naturforschung* 2B:425-427.
- GUILLEMAT, J., and J. LAMBERT. 1960. I. Etude du pouvoir fungicide du carbamate. II. Etude des mélanges carbamate et sels cupriques. *Phytopathologie-Phytopharmacie* 9:7-21.
- HAMILTON, JAMES M. 1959. Evaluation of fungicides in the greenhouse, p. 253-257. In C. S. Holton [ed.] *Plant Pathology, Problems and Progress, 1908-1958*. The University of Wisconsin Press, Madison.
- HENRY, BERT W., and ELIZABETH C. WAGNER. 1940. A rapid method of testing the effects of fungicides on fungi in culture. *Phytopathology* 30:1047-1049.

- HEUBERGER, JOHN W. 1940. A laboratory biological assay of tenacity of fungicides. *Phytopathology* 30:840-847.
- HEYNS, A. J., G. A. CARTER, K. ROTHWELL, and R. L. WAIN. 1965. Investigations on fungicides. XII. The fungicidal activity of certain N-carboxymethylthiocarbamic acid derivatives. *Annals of Applied Biology* 56:399-409.
- HIMELICK, E. B., and DAN NEELY. 1965. Bioassay using cellophane to detect fungistatic activity of compounds translocated through the vascular system of trees. *Plant Disease Reporter* 49:949-953.
- HISLOP, E. C., and P. O. PARK. 1962. Studies on the chemical control of *Phytophthora palmivora* (Butl.) Butl. on *Theobroma cacao* L. in Nigeria. I. Laboratory bio-assay of fungicides on detached pods. *Annals of Applied Biology* 50:57-65.
- HOLLOMAN, ARTHUR, JR., and ROY A. YOUNG. 1951. Selection by laboratory assay of fungicides for field control of *Botrytis* leaf spot of gladiolus. *Phytopathology* 41:940. (Abstr.)
- HORN, NORMAN L. 1964. A detached cucumber leaf method of screening fungicides for the control of anthracnose. *Plant Disease Reporter* 48:331-332.
- HORSFALL, JAMES G. 1945a. Fungicides and their action. *Chronica Botanica Company*, Waltham, Massachusetts. 239 p.
- . 1945b. Quantitative bioassay of fungicides in the laboratory. *Botanical Review* 11:357-397.
- . 1951. Relative specificity of *Sclerotinia* and *Stemphylium* to toxicants. *Phytopathology* 41:658. (Abstr.)
- . 1956. Principles of fungicidal action. *Chronica Botanica Company*, Waltham, Massachusetts. 279 p.
- , J. W. HEUBERGER, and A. E. DIMOND. 1941. Predicting protective value of fungicides in the laboratory. *Phytopathology* 31:12. (Abstr.)
- , ———, E. G. SHARVELLE, and J. M. HAMILTON. 1940. A design for laboratory assay of fungicides. *Phytopathology* 30:545-563.
- , and RAYMOND J. LUKENS. 1966. Selectivity of fungicides. *Connecticut Agricultural Experiment Station Bulletin* 676. 24 p.
- HOWARD, FRANK L. 1939. The values of testing fungicides in the laboratory before use in the field. *Proceedings of the American Society for Horticultural Science* 37:409-414.
- KEIL, HARRY L., JAMES G. HORSFALL, and SAUL RICH. 1952. Relation of laboratory ED50 to field performances of fungicides. *Phytopathology* 42:113. (Abstr.)
- KIRBY, A. H. M., and E. L. FRICK. 1953. Modifications in the glass-slide spore-germination test for use in fungicide research. *Annual Report for 1952, East Malling Research Station (Kent, England)* 36:161-167.
- KLOMPARENS, WILLIAM, and JOHN R. VAUGHN. 1952. The correlation of laboratory screening of turf fungicides with field results. *Michigan Agricultural Experiment Station Quarterly Bulletin* 34:425-435.
- KOOPMANS, M. J. 1959. An in vitro evaluation of the toxicity of chemicals for Erysiphaceae. *Mededelingen van de Landbouwhogeschool en de opzoekingsstations van de staat te Gent* 24:821-827.
- KOVACS, ANDREA. 1961. Ricerche sulla persistenza dei fungicidi organici. I. L'influenza del tipo di superficie sulla dilavabilità. *La Ricerca Scientifica, Parte II: Rendiconti, Sezione B: Biologica* 1:57-66.
- KUIFUS, K. H. 1957. Beitrag zur methodik der Fungizidprüfung von Nass- und Trockenbeizmitteln. *Phytopathologische Zeitschrift* 28:281-284.
- KUNDERT, J. 1956. Über die Anwendung statistischer methoden bei der biologischen Prüfung von Fungiziden. *Annuaire agricole de la Suisse. N.S.* 5, 6, 70:687-709.
- LEBEN, CURT. 1949. The determination of antimycin on plant leaves. *Phytopathology* 39:13. (Abstr.)
- , and G. W. KERT. 1949. Laboratory and greenhouse studies of antimycin preparations as protectant fungicides. *Phytopathology* 39:529-540.
- , and ———. 1950. A bioassay for tetramethylthiuramdisulfide. *Phytopathology* 40:950-954.
- LEE, H. ATHERTON, and J. P. MARTIN. 1927. A method for testing in vitro the toxicity of dust fungicides to fungous spores. *Phytopathology* 17:315-319.
- LE TOURNEAU, DUANE, and LAWRENCE BUER. 1961. The toxicity of some chlorinated phenols and aryloxyalkanecarboxylic acids to *Verticillium albo-atrum*. *Phytopathology* 51:128-129.
- LITCHFIELD, J. T., JR., and F. WILCOXON. 1949. A simplified method of evaluating dose-effect experiments. *Journal of Pharmacology and Experimental Therapeutics* 96:99-113.
- LUIJTEN, J. G. A., and G. J. M. VAN DER KERK. 1961. Investigations on organo-tin compounds. XIV. Preparation and antifungal properties of some triaryl- and triaralkyl-tin acetates. *Journal of Applied Chemistry* 11:35-37.
- MANOELS, G. R., and R. T. DARBY. 1953. A rapid cell volume assay for fungitoxicity using fungus spores. *Journal of Bacteriology* 65:16-26.
- , and R. C. H. SIU. 1950. Rapid assay for growth: Determination of microbiological susceptibility and fungistatic activity. *Journal of Bacteriology* 60:249-262.
- MANTEN, A., H. L. KLÖPPING, and G. J. M. VAN DER KERK. 1950. Investigations on organic fungicides. II. A new method for evaluating antifungal substances in the laboratory. *Antonie van Leeuwenhoek Journal of Microbiology and Serology* 16:282-294.
- MARSH, R. W. 1936. Notes on a technique for the laboratory evaluation of protective fungicides. *British Mycological Society Transactions* 20:304-309.
- MARTIN, HUBERT. 1942. The significance of the bio-assay in studies of fungicidal action. *Annals of Applied Biology* 29:326-328.
- MASON, CURTIS L., and DWIGHT POWELL. 1947. A Pythium plate method of evaluating fungicides. *Phytopathology* 37:527-528.
- MCCALLAN, S. E. A. 1930. Studies on fungicides. II. Testing protective fungicides in the laboratory. *New York (Cornell) Agricultural Experiment Station Memoir* 128:8-24.
- . 1947. Bioassay of agricultural fungicides. *Agricultural Chemicals* 2(9):31-34, 67; 2(10):45.
- . 1948. Evaluation of chemicals as seed protectants by greenhouse tests with peas and other seeds. *Contributions from Boyce Thompson Institute* 15:91-117.
- . 1959. Evaluation of fungicides in the laboratory, p. 248-253. *In* C. S. Holton [ed.] *Plant Pathology, Problems and Progress, 1908-1958*. The University of Wisconsin Press, Madison.
- . 1966. Evaluation of fungicides in the greenhouse, p. 191-215. *In* Symposium on Plant Diseases in the Pacific, Pacific Science Congress, 11th, Tokyo, August 25-27, 1966.
- , H. P. BUCHFIELD, and L. P. MILLER. 1959. Interpretation of dosage-response curves. *Phytopathology* 49:544-545. (Abstr.)
- , and LAWRENCE P. MILLER. 1957. Effect of fungicides on oxygen consumption and viability of mycelial pellets. *Contributions from Boyce Thompson Institute* 18:483-495.
- , and ———. 1958. Innate toxicity of fungicides. *Advances in Pest Control Research* 2:107-134.
- , ———, and RICHARD M. WEED. 1954. Comparative effect of fungicides on oxygen uptake and germination of spores. *Contributions from Boyce Thompson Institute* 18:39-68.
- , and R. H. WELLMAN. 1942. Fungicidal versus fungistatic. *Contributions from Boyce Thompson Institute* 12:451-463.

- , —, and FRANK WILCOXON. 1941a. The different toxicity rating of compounds by different fungi and experiments in spore-germination tests of fungicides. *Phytopathology* 31:16. (Abstr.)
- , —, and —. 1941b. An analysis of factors causing variation in spore germination tests of fungicides. III. Slope of toxicity curves, replicate tests, and fungi. *Contributions from Boyce Thompson Institute* 12:49-77.
- , and FRANK WILCOXON. 1938. Laboratory comparisons of copper fungicides. *Contributions from Boyce Thompson Institute* 9:249-263.
- , and —. 1939. An analysis of factors causing variation in spore germination tests of fungicides. I. Methods of obtaining spores. *Contributions from Boyce Thompson Institute* 11:5-20.
- MCINTOSH, A. H. 1961. Some variants and possible errors in the test-tube dilution and slide-germination methods for laboratory testing of fungicides. *Annals of Applied Biology* 49:424-432.
- MILDNER, P., BRANKA MIHANOVIC, MIRNA JUSIC, M. HAJSIG, and V. KUZMANOVIC. 1963. Antifungal action of alkoxy ethyl mercuric acetates. *Antonie van Leeuwenhoek Journal of Microbiology and Serology* 29:421-424.
- MILLER, HAROLD J. 1943. A comparison of laboratory and field retention and protective value of certain copper fungicides. *Phytopathology* 33:899-909.
- , 1949. Modifications of the slide-germination method of evaluating fungicides including the use of *Venturia inaequalis* and *Phytophthora infestans*. *Phytopathology* 39:245-259.
- MILLER, LAWRENCE P., S. E. A. MCCALLAN, and RICHARD M. WEED. 1953. Rate of uptake and toxic dose on a spore weight basis of various fungicides. *Contributions from Boyce Thompson Institute* 17:173-195.
- MISATO, T. 1963. Correlation between the fungicidal activity in vitro and the protectant effect in vivo, p. 32. In *Abstracts, Fifth International Pesticides Congress, London, July 17-23, 1963*. International Union Pure & Applied Chemistry, London.
- MONROE, CARL MAX. 1963. Comparison of in vitro and in vivo bioassay of foliage fungicides. *Dissertation Abstracts* 24(3):926.
- MONTGOMERY, H. B. S., and M. H. MOORE. 1938. A laboratory method for testing the toxicity of protective fungicides. *Journal of Pomology and Horticultural Science* 15:253-266.
- MOREAU, CLAUDE, and MIREILLE MOREAU. 1959. Technique simple de comparaison du pouvoir fongicide de diverses substances. *Revue de Mycologie* 24:59-64.
- NEELY, DAN, and E. B. HIMELICK. 1966. Simultaneous determination of fungistatic and fungicidal properties of chemicals. *Phytopathology* 56:203-209.
- NIEMANN, G. J., and J. DEKKER. 1966. Evaluation of the activity of a number of procaine derivatives and analogous compounds against powdery mildew (*Sphaerotheca fuliginea*) on cucumber. *Annals of Applied Biology* 57:53-58.
- NISIKADO, Y., T. NAKAYAMA, and Y. MIYAWAKI. 1951. Studies on the methods for testing the effectiveness of fungicides. I. The use of slide glasses as a medium for testing spray materials. *Berichte des Ohara Instituts für Landwirtschaftliche Forschungen (Okayama, Japan)* 9:317-328.
- PALMITER, D. H., and G. W. KEITT. 1937. The toxicity of copper-lime-arsenic mixtures to certain phytopathogenic fungi grown on malt agar plates. *Journal of Agricultural Research* 55:439-451.
- PETERSON, P. D. 1941. The spore-germination method of evaluating fungicides. *Phytopathology* 31:1108-1116.
- PIANKA, M., J. D. EDWARDS, and C. B. F. SMITH. 1966. Studies in fungitoxicity. V. Fungicidal activity of certain dithiocarbamates. *Journal of the Science of Food and Agriculture* 17:407-416.
- PLUIJGERS, C. W., and A. KAARS SJPESTEIJN. 1966. The antifungal activity of thiosemicarbazides in vitro and their systematic activity against cucumber scab *Cladosporium cucumerinum*. *Annals of Applied Biology* 57:465-473.
- POWELL, DWIGHT. 1961. A "water-bridge" technique for evaluating fungicides. *Illinois Research* 3(1):16.
- PRUSOVA, HELENA. 1962. Metoda vysusenych kapek fungicidni suspence. *Ceska Mykologie* 16:214-218.
- REDDICK, DONALD, and ERRETT WALLACE. 1910. On a laboratory method of determining the fungicidal value of a spray mixture or solution. *Science* 31:798.
- RICH, SAUL, JAMES G. HORSFALL, and HARRY L. KEIL. 1953. The relation of laboratory to field performance of fungicides, p. 282-287. In *Report of the Thirteenth International Horticultural Congress, London, 1952*. Royal Horticultural Society, London.
- SCHMIDT, H. 1951. Laborshnelltest zur Fungizidprüfung. *Nachrichtenblatt des Deutschen Pflanzenschutzdienstes, Berlin*. N. F. 5, 11:208-212.
- SERRA, G. 1964. Etude de la dégradation du captane du phaltane et du difolatan sous l'influence de la lumière. *Phytiatrie-Phytopharmacie* 13:107-110.
- SHAFFER, WILLIAM B. 1952. Fungitoxicity investigations using a novel spore germination technique. *Phytopathology* 42:519. (Abstr.)
- SHARVELLE, E. G., and E. N. PELLETIER. 1956. Modified paper-disk method for laboratory fungicidal bioassay. *Phytopathology* 46:26. (Abstr.)
- SMITH, R. J., and W. H. READ. 1961. Investigations on fungitoxic derivatives of salicylaldehyde. 1. The 3-, 5- and 3, 5-chlorinated derivatives of salicylanilide and salicyl-*para*-chloranilide. *Annals of Applied Biology* 49:102-109.
- SPENCER, D. M. 1962. Polystyrene microbeakers for spore germination tests. *Plant Pathology* 11:41-43.
- TAMURA, HIROKUNI. 1954. Study on the evaluating method of organic fungicides. On the "petri dish method" as a new method of bioassay in laboratory. (In Japanese. English resume.) *Bulletin of the National Institute of Agricultural Sciences, Series C*, 4:95-107.
- THORNBERRY, H. H. 1950. A paper-disk method for the quantitative evaluation of fungicides and bactericides. *Phytopathology* 40:419-429.
- TORGESON, D. C. 1967. Determination and measurement of fungitoxicity, p. 93-123. In *Dewayne C. Torgeson [ed.] Fungicides, an advanced treatise, Vol. I*. Academic Press, Inc., New York.
- WAGGONER, PAUL E., SAUL RICH, and J. G. HORSFALL. 1952. Prediction of field performance of fungicides. *Phytopathology* 42:477. (Abstr.)
- WALKER, ALMA TOEVS. 1955. Germination and respiration responses of *Myrothecium verrucaria* to organic fungicides. *Iowa State College Journal of Science* 30:229-241.
- WELLMAN, R. H. 1967. Commercial fungicide development, p. 125-151. In *Dewayne C. Torgeson [ed.] Fungicides, an advanced treatise, Vol. I*. Academic Press, Inc., New York.
- , and F. D. HEALD. 1940. The toxicity of certain chemicals in aqueous solutions to spores of *Penicillium expansum*. *Phytopathology* 30:638-648.
- , and S. E. A. MCCALLAN. 1943. Correlations within and between laboratory slide-germination, greenhouse tomato foliage disease, and wheat smut methods of testing fungicides. *Contributions from Boyce Thompson Institute* 13:143-169.
- WILCOXON, FRANK, and S. E. A. MCCALLAN. 1939. Theoretical principles underlying laboratory toxicity tests of fungicides. *Contributions from Boyce Thompson Institute* 10:329-335.
- YOUNG, J. E. 1944. A simplified laboratory method of testing the toxicity of fungicides. *Illinois Academy of Science Transactions* 37:59-62.

