

Q.630.7

146c


no.907

1983

cop.5



UNIVERSITY OF  
ILLINOIS LIBRARY  
AT URBANA-CHAMPAIGN  
AGRICULTURE



Digitized by the Internet Archive  
in 2012 with funding from  
University of Illinois Urbana-Champaign

# Weed Management Guide 1983

## FOR COMMERCIAL VEGETABLE GROWERS

---

*Restricted-use herbicides are identified with an asterisk(\*)*.

*You must be certified as a pesticide applicator to use restricted-use pesticides.*

*See your county Extension adviser in agriculture for information.*

WEED GROWTH reduces the income of vegetable growers in the United States by millions of dollars annually as a result of lower yields, poorer quality, and added labor in harvesting and processing.

This guide should be used together with the grower's knowledge of soil types and the crop and weed history of the area to be treated. Whether to use herbicides or other means of weed control depends in part on the severity of past weed infestations. In some instances, mechanical control may be sufficient. If so, shave off weeds with a sharp hoe or cultivator while gently breaking up the crust. Deep tillage causes severe injury to many shallow-rooted plants and helps place a fresh supply of weed seeds in position to germinate. Keeping equipment sharp and in good condition will help reduce injury to desirable plants. Hoe carefully around your plants, and hand pull weeds close to the plants.

For warm season crops such as fresh market tomatoes, peppers, eggplant, okra, cucumbers, and melon, black polyethylene mulch will control annual weeds, conserve moisture, and increase the soil temperature in early spring. The higher temperature increases early season growth. Natural mulch materials may require considerable hand labor for application. Most organic materials are bulky and must be hauled to the place of use. This is a problem for large commercial plantings. Organic mulches tend to reduce soil temperature.

Herbicide application may be needed in addition to mechanical control. Several herbicides may be suggested for some crops. These herbicides have shown good control with no injury to the vegetables under test conditions. Not all herbicides cleared for use on a species are necessarily listed. Where the choice of more than one herbicide is suggested, the decision rests with the grower and is based on his knowledge of past weed infestations, crop rotations, and material costs. Where one herbicide will not control the weeds present, a combination of herbicides may be suggested. When using a herbicide for the first time, it is advisable to use a small-scale trial.

These suggestions for weed control in vegetables are based on research at the Illinois Agricultural Experiment Station, the U.S. Department of Agriculture, and other research institutions. The University of Illinois and its agents assume no responsibility for results from the use of herbicides, whether or not they were used in accordance with suggestions, recommendations, or directions of the manufacturer or any governmental agency.

*Reading the label of the herbicide container is the most profitable time you spend in weed control. Use of the material and methods of application and use depend on registration of the herbicide by federal and state Environmental Protection Agencies (EPA). Do not use any herbicide unless the label states that it is cleared for the use on the crop to be treated.*

Herbicides are being classified for *general use* or *restricted use* by the U.S. Environmental Protection Agency. A person wishing to use a herbicide classified for restricted use must be certified as a private or commercial pesticide applicator by the Illinois Department of Agriculture. Contact your county Extension adviser in agriculture for details about this program.

Only a few herbicides have been classified at this time. More may be classified later.

When applying mixtures of chemicals, the *user* assumes responsibility for freedom from residues if the mixture is not labeled by the EPA.

Suggestions sometimes change during the growing season, based on EPA clearances that were made after this circular was issued. This publication, printed once a year, is subject to change without notification.

Watch for notice of changes in the EPA registration of herbicides (as released by the EPA) in the *Illinois Vegetable Farmer's Letter* and the *Insect, Weed, and Plant Disease Survey Bulletin*. Subscription forms for the latter are available from the Agricultural Newsletter Service, 116 Mumford Hall, 1301 West Gregory Drive, Urbana, Illinois 61801, or your county Extension Office. You can obtain the *Vegetable Farmer's Letter* from Vegetable Crops Extension, University of Illinois, 1103 West Dorner Drive, Urbana, Illinois 61801.

## For Application During the Growing Season (1983 Only)

Crop	Treatment	Active ingredient per acre actually covered <sup>a</sup>	Weeds controlled	Timing of application (based on crop stage)	Remarks, cautions, limitations
Asparagus (seedlings)	Amiben	3 lb.	Annuals	Immediately after seeding	Irrigation or rainfall after treatment will give maximum control.
Asparagus (established plantings) <sup>b,c</sup>	dalapon	5-10 lb.	Perennial grass	End of harvest season following disking	Apply when grass weeds are 3 to 4 in. tall. Direct spray under fern growth. Use surfactant as directed on label.
	diuron	1-4 lb.	Annuals	In spring, after harvest, or both	Apply after disking. Do not exceed 6 lb. per growing season; use a lighter rate on sandy soil. With diuron and Princep, a spring application may be sufficient after the first year.
	Princep	3-4 lb.	Annuals	In spring, after harvest, or both	Apply after disking. Do not treat during the last year in asparagus because of residue.
	Sinbar	1.2-2.4 lb.	Annuals	In spring, after harvest, or both	Use lower rates on coarse soils. Do not apply more than 2.4 lb. per acre per year. Do not use on soils with less than 1 percent organic matter. Do not plant to any other crop for two years after application.
	metribuzin	1-2 lb.	Primarily broad- leaf weeds	Early spring before the spears emerge or after harvest	Apply after disking. Do not apply within 14 days of harvest. Can help control broadleaf weeds when used with dalapon, diuron, or Princep. Do not apply more than 2 lb./acre per growing season.
Beans, dry, lima and snap <sup>d</sup>	<b>Preemergence</b>				
	Treflan	0.5-0.75 lb.	Annuals* (primarily grasses)	Preplant soil application, in- corporate with soil immediately	Plant crop immediately, or within 3 weeks after applica- tion. Can use up to 1 lb. per acre on dry beans.
	Basalin	0.75-1.5 lb.	Annuals (primarily grasses)	Preplant soil application, incor- porate with soil immediately	
	dinoseb	6-7.5 lb.	Annuals	Can be used between planting and crop emergence	Do not use on light, sandy soil. Some stand reduction may result from use. See label for precautions.
	<b>Postemergence</b>				
	Basagran	0.75-1 lb.	Annual broad- leaf weeds, Canada thistle, nutsedge	When weeds are small and are actively growing; after the first trifoliolate leaf appears on beans	Can provide good, broad-spectrum control when combined with a grass-active herbicide. Do not mix with other pes- ticides. See Basagran entry under corn, postemergence for Canada thistle and nutgrass control.
Beans, snap	Eptam	3 lb.	Annual grasses and nutgrass <sup>f</sup>	Preplant soil application, incor- porate with soil immediately	
	Eptam +	2-3 lb.		Preplant soil application, incor- porate with soil immediately	TRIAL USE IN 1983. Research results have shown this combination to control a broader spectrum of weeds than either herbicide alone.
	Treflan	0.5-0.75 lb.			
	Dacthal	6-10 lb.	Annuals* (primarily grasses)	Immediately after seeding	Do not feed treated plant parts to livestock.
Beans, lima and dry	Amiben	2-3 lb.	Broad spectrum of annual weeds	Immediately after seeding, or preplant-incorporated for lima beans	Field may be rotary-hoed without destroying herbicide action.
	Dual	1.5-3 lb.	Annuals	Preplant soil application, incor- porate with soil, or pre- emergence	
	Lasso	2-3 lb.	Annuals	Lima beans: preplant soil ap- plication, incorporate into upper 1 to 2 inches. Dry beans: preplant soil application, incor- porate into upper 1 to 2 inches, or preemergence	
Beans, dry	Eptam +	2-3 lb.		Preplant soil application, incor- porate with soil immediately	TRIAL USE IN 1983. Research results have shown this combination to control a broader spectrum of weeds than either herbicide alone.
	Treflan	0.5-0.75 lb.			
Beets, garden <sup>d</sup>	Pyramin	4 lb.	Annuals (primarily broadleaved)	Preemergence or after beets emerge and before weeds have two true leaves	Rainfall or irrigation needed to activate. Where grasses are a severe problem, use 4 lb. of Pyramin plus 4 lb. of Ro-Neet.
	Ro-Neet	4 lb.	Annual grasses	Preplant soil application, incor- porate with soil immediately	Use a combination treatment with Pyramin to broaden control spectrum.
Broccoli <sup>d</sup> Brussels sprouts <sup>d</sup> Cabbage <sup>d</sup> Cauliflower <sup>d</sup>	<b>Direct-seeded or transplanted</b>				
	Treflan	0.5-0.75 lb.	Annuals* (primarily grasses)	Preplant soil application, incor- porate with soil immediately	Stunting or growth reduction may occur at recom- mended rates under growth stress conditions. Can use up to 1 lb. per acre on transplants.
	Dacthal	6-10 lb.	Annuals* (primarily grasses)	Immediately after seeding. Can also be incorporated preplant	
Carrots <sup>d</sup>	<b>Preemergence</b> Treflan	0.5-1 lb.	Annuals* (primarily grasses)	Preplant soil application, incorporate with soil immedi- ately	Seed after application to 3 weeks later.

Carrots continued on the next page.

All notes are at the end of this table (page 6).



## For Application During the Growing Season (continued)

Crop	Treatment	Active ingredient per acre actually covered <sup>a</sup>	Weeds controlled	Timing of application (based on crop stage)	Remarks, cautions, limitations
Cucumbers Muskmelons <sup>c</sup> Watermelons <sup>c</sup>	Alanap L	3-5 lb.	Annuals <sup>f</sup>	Immediately after seeding or transplanting	Do not use on cold soil. Rainfall or irrigation after treatment gives maximum control.
	Prefar	4-6 lb.	Annuals (primarily grasses)	Preplant soil application, incorporate with soil immediately	Keep away from foliage. Apply to soil after the weeds have been removed. Is primarily a grasskiller. Consult label for sensitive crops within 18 months after application. Can be used in rotation with tomatoes, broccoli, cauliflower, lettuce, carrots, onions, and summer squash within 18 months. Soybeans can be planted 12 months after application.
	Prefar plus Alanap L	4 lb. +2-3 lb.	Grasses and broadleaved weeds	Preplant light incorporation	Has value for broad-spectrum weed control. Consult label for sensitive crops within 18 months after Prefar application. Has EPA approval as a tank mixture.
Where earliness is desired, black polyethylene mulch can be used as an alternative to herbicides. It will control annual weeds, conserve moisture, and increase the soil temperature in early spring.					
Eggplant	Dacthal	6-10 lb.	Annuals <sup>g</sup> (primarily grasses)	After plants are established, 4-6 weeks after transplanting	Cultivate and weed prior to application. Can be applied to plants as part of a uniform soil application.
	Devrinol	1-2 lb.	Annuals	Preplant soil incorporation	For use in transplanted eggplant.
Where earliness is desired, black polyethylene mulch can be used as an alternative to herbicides. It will control annual weeds, conserve moisture, and increase the soil temperature in early spring.					
Greens (for beets, mustard greens, and spinach — see note d)	Dacthal	6-10 lb.	Annuals <sup>g</sup> (primarily grasses)	Immediately after seeding	For use on collards, kale, mustard greens, and turnips.
	Treflan	0.5-0.75 lb.	Annuals <sup>g</sup> (primarily grasses)	Preplant soil application, in- corporate with soil immediately	For use on collards, kale, mustard greens, and turnip greens.
	Furloe	1-2 lb.	Primarily broad- leaved annuals	Preemergence	For spinach only. Use lower rates in cool, wet weather.
Horseradish <sup>d</sup>	Dacthal	6-10 lb.	Annuals <sup>g</sup> (primarily grasses)	Immediately after transplanting	
Lettuce <sup>e,d</sup>	Balan	1.5 lb.	Annuals	Preplant soil incorporation Incorporate with soil immediately	Is primarily a grasskiller. Seed after application to 3 weeks later. Do not plant wheat, barley, rye, grass, onions, oats, beets, or spinach for 12 months after application.
	Kerb <sup>h</sup>	1-2 lb.	Annuals	Preemergence or preplant- incorporated	Do not use when the air temperature exceeds 85° F. Use the lower rates listed on sandy soil. Do not use on peat or muck soils. See label for rotation crops. For best results, rainfall or irrigation is needed 1 to 2 days after application, especially during warm weather.
Okra <sup>d</sup>	Dual	1.5-3 lb.	Annuals	Preplant soil application, incorporate with soil, or preemergence	
	Enide	3-5 lb.	Annuals	At planting	
	Treflan	0.5-1 lb.	Annuals (primarily grasses)	Preplant soil application, incorporate with soil immediately	
Where earliness is desired, black polyethylene mulch can be used as an alternative to herbicides. It will control annual weeds, conserve moisture, and increase the soil temperature in early spring.					
Onions <sup>b</sup>	<b>Preemergence</b>				
	Dacthal	6-10 lb.	Annuals <sup>g</sup> (primarily grasses)	Immediately after seeding or transplanting	May not kill smartweed or common ragweed. Can be used on seeds, sets, or seedlings. Use only on mineral soils. Use lower rates on sandy soils. A double application of Dacthal can be used at seeding, layby, or both.
	Randox	4-6 lb.	Annuals <sup>i</sup> (primarily grasses)	Just before onions emerge	Use on muck soils. Heavy rainfall may reduce stand. Very effective on purslane and pigweed.
	<b>Postemergence</b>				
Furloe	3-6 lb.	Broadleaved weeds (especially smartweed)	On seeded onions: loop stage or after 3- to 4-leaf stage	In later sprays, direct at base of onion plant. If applied more than once, do not exceed 6 lb. per acre for the season. Use lower rates in cool, wet weather. Use no later than 30 days before harvest. Do not use on sandy soils.	
Brominal	0.25-0.38 lb.	Broadleaved weeds	When onions have 2 to 5 true leaves	TRIAL USE IN 1983. Use 50 to 70 gallons of water per acre. Apply when onion foliage is dry for greatest crop safety. Suggested temperature for spraying is 80° F with low humidity.	
Peas <sup>d</sup>	<b>Preemergence</b>				
	propachlor	4-4.9 lb.	Annuals	Preemergence	Do not use on sandy soil.
Treflan	0.5-0.75 lb.	Annuals <sup>g</sup>	Preplant soil incorporation, incorporate with soil immediately	Seed after application to 3 weeks later. Some reduction of growth and stand reduction possible under stress. May suppress some root rot.	

Peas continued on the next page.

All notes are at the end of this table (page 6).



## For Application During the Growing Season (continued)

Crop	Treatment	Active ingredient per acre actually covered <sup>a</sup>	Weeds controlled	Timing of application (based on crop stage)	Remarks, cautions, limitations
Peas (continued) <sup>d</sup>	Treflan + Surflan	0.5 lb. + 0.5 lb.	Annuals	Preplant soil application, incorporate with soil immediately	Do not use on soils of less than 1.5 percent organic matter. May suppress <i>Aphanomyces</i> root rot. May broaden weed control more than either herbicide alone. Do not feed forage to livestock. Do not plant any root crop for 12 months after application.
	Basalin	0.75-1.5 lb.	Annuals (primarily grasses)	Preplant soil application, incorporate with soil immediately	
	Dual	1.5-3 lb.	Annuals	Preplant soil application, incorporate with soil, or preemergence	
	<b>Preemergence or Postemergence</b> dinoseb	0.3-9 lb.	Annuals (primarily broad- leaved weeds)	Preemergence or postemergence	Preemergence, use 6 to 9 lb.; postemergence, use 0.3 to 1.1 lb. Apply prior to bloom when peas are 2 to 8 in. tall. See label for further precautions. Preemergence use may help suppress root rot.
	<b>Postemergence</b> Basagran	0.75-1 lb.	Annual broad- leaved weeds, Canada thistle, nutsedge	When weeds are small and are actively growing; after peas have 3 pairs of leaves (or 4 nodes)	Can help control Canada thistle. Can provide good, broad- spectrum control when used with a grass-active herbicide. Do not mix with other pesticides. See Basagran entry under corn, postemergence for Canada thistle and nut- grass control.
	MCPB	1 lb.	Broadleaved weeds and Canada thistle	When peas are 3-7 in. tall and no later than 4 nodes prior to pea blossom	May delay maturity 1 to 4 days. Use at least 20 gallons of water per acre. Do not feed vines to livestock. MCPA is more effective on mustard. MCPB or Vacate may be less injurious to peas.
	MCPA (Na salt)	0.25-0.5 lb.			
Vacate (MCPA amine)	0.115-0.154 lb.				
Potatoes, Irish <sup>e,d</sup>	Dual	1.5-3 lb.	Annuals	Drag-off treatment at emergence	
	Eptam	3-6 lb.	Annual grasses and nutgrass <sup>f</sup>	Drag-off treatment at emer- gence or preplant soil application; incorporate with soil immediately	Use lower rate on sandy soil.
	Treflan	0.5-1 lb.	Annuals* (primarily grasses)	Drag-off treatment at emergence	Use a light incorporation.
	Lorox	0.75-2 lb.	Annuals	Apply prior to potato emergence	Plant tubers at least 2 in. deep. Do not replant treated area to other crops for 4 months after treatment. May injure crop on light, sandy soil. Do not apply over exposed tubers.
	dalapon	7 lb.	Quackgrass	Before plowing in spring; wait 4 days before plowing and planting	Not for fields intended for red-skinned varieties or White Rose. Do not plant potatoes for 4 weeks. Use surfactant as directed on label.
	metribuzin	0.25-0.5 lb.	Annuals (primarily broadleaved)	Postemergence, following a preemergence grass herbicide	Can be used preemergence also. Do not exceed 1 lb. per acre in a season. Do not apply within 60 days of harvest. Do not use on red-skinned or early-maturing white varieties. Do not apply in cool, wet weather.
	Lasso	2.5-3 lb.	Annuals	Apply at drag-off	Do not use on sandy soils. Can be used alone or in com- bination with Lorox or dinoseb.
Potatoes, sweet <sup>b</sup>	Dacthal	6-10 lb.	Annuals* (primarily grasses)	Immediately after planting	
	Amiben	3 lb.	Annuals	Immediately after planting	
	Enide	4-6 lb.	Annuals	Immediately after trans- planting	Do not plant nonapproved crops on treated soil during the same season.
Squash Pumpkins	Amiben	3-4 lb.	Annuals	As soon after seeding as possible, or preplant- incorporated	Use on loam soils. Amiben can be applied broadcast or banded over the row in pumpkins.
	Prefar	4-6 lb.	Annuals (primarily grasses)	Preplant soil application, incorporate with soil immediately	Use on sandy soils. Is primarily a grasskiller. Consult label for sensitive crops within 18 months after applica- tion. Can be used in rotation only with tomatoes, broccoli, cauliflower, lettuce, carrots, onions, and summer squash within 18 months of application. Soybeans can be planted 12 months after application.
Pumpkins	propachlor	4-5 lb.	Annuals	Preemergence	Do not use on sandy soils.
Tomatoes, direct-seeded and trans- planted <sup>c</sup>	Enide	4-6 lb.	Annuals	Preemergence	Do not plant other food crops on treated areas for 6 months. If used under dry soil conditions, a shallow (1 in.) incorporation as a preplant treatment may improve weed control. Can also be used on transplanted peppers.
	Devrinol	1-2 lb.	Annuals	Preplant soil incorporated	Also used on direct-seeded and transplanted peppers. Enide + Devrinol is labeled as a tank mixture.

Tomatoes continued on the next page.

All notes are at the end of this table (page 6).

## For Application During the Growing Season (continued)

Crop	Treatment	Active ingredient per acre actually covered <sup>a</sup>	Weeds controlled	Timing of application (based on crop stage)	Remarks, cautions, limitations
Tomatoes (continued) <sup>c</sup>	metribuzin	0.25-1 lb. (min.-max.)	Primarily broad-leaf. Should be used with a grass-active herbicide.	Preplant incorporated. Post-emergence, can be broadcast or directed.	Apply with ground equipment to seeded and transplanted tomatoes. Do not use air-blast or other high-pressure spray equipment. Do not use on peppers.
		0.25-0.5 lb.		Preplant incorporated, transplant tomatoes	Alone or in a tank-mix combination with Treflan.
		0.25-0.5 lb.		Broadcast spray, established tomatoes	Single or multiple applications. Minimum of 14 days between treatments. Direct-seeded plants should have 5 or 6 leaves; transplants should show new growth.
		0.5-1 lb. (For min.-max. rates)		Directed spray, established tomatoes	Recommended for use in fields with severe weed problems, or for fields with hard-to-control weeds. Do not apply within 7 days of harvest, or within 3 days following cool, wet, or cloudy weather; otherwise, crop injury may occur. Do not apply to established tomatoes within 24 hours after application of other pesticides. Do not apply more than 1 lb./acre per crop season, or more than 1 lb./acre within a 35-day period. Allow at least 14 days between applications, regardless of the dosage or method used. Do not use hot caps on tomatoes within 7 days before application, or at any time afterward. Do not tank-mix with other pesticides, except Treflan.
Tomatoes and Peppers, transplanted	Amiben	3-4 lb.	Annuals	Wait 3 days after transplanting to apply	Use granular formulation only. Apply to dry foliage in order to avoid leaf burn. Do not use on sandy soils.
	Treflan	0.5-1 lb.	Annuals <sup>e</sup> (primarily grasses)	Preplant soil application, incorporate with soil immediately	Some reduction of growth may be possible under growth stress conditions, or if rates are higher than suggested for the soil type.

Where earliness is desired, black polyethylene mulch can be used as an alternative to herbicides. It will control annual weeds, conserve moisture, and increase the soil temperature in early spring.

**\* Restricted-use herbicide.**

<sup>a</sup> Based on active ingredients (actual amount of active herbicide in material or acid equivalent). Use lower rate on sandy soil and higher rate on clay and loam soils. When using a band application over the row, adjust amount of material applied to the part of an acre treated. <sup>b</sup> For perennial weed control, applications during and outside the growing season, see the following table. <sup>c</sup> For stale seedbeds, before crop emergence, see the following table. <sup>d</sup> For perennial grass control, applications outside the growing season, see the following table. <sup>e</sup> May not control ragweed and panicum. <sup>f</sup> May not control smartweed. <sup>g</sup> May not control ragweed, smartweed, and velvetleaf. <sup>h</sup> May not control crabgrass. <sup>i</sup> May not control smartweed and velvetleaf.

## For Application Outside the Growing Season

Asparagus	Stale seedbed, before crop emergence				
	paraquat*	0.5-1 lb.	All emerged green foliage	Before crop emergence; allow maximum weed emergence prior to treatment	Weeds that emerge after treatment will not be controlled. Crop plants that have emerged at application will be injured. Do not apply within 18 months of harvest. Use with a preemergence or preplant sustained-action weed control system.
Corn, sweet Lettuce Melons Peppers Potatoes Tomatoes	paraquat*	0.5-1 lb.	All emerged green foliage	Before crop emergence	Weeds that emerge after treatment will not be controlled. Crop plants that have emerged at application will be injured. Use with a preemergence or preplant, sustained-action weed control system.
Asparagus	Perennial weed control, applications during and outside the growing season				
	Roundup	2-5 lb.	(See remarks)	Before emergence, or with shielded or directed sprays during fern growth	Use to control milkweed, thistle, field bindweed, quackgrass, or Johnsongrass. Apply to quackgrass when it is 6-8 in. tall in the fall or spring. Apply to Johnsongrass when it is at least 12 in. tall and actively growing. Do not till for the specified time for each species (see label). Does not provide residual weed control. Do not mix, store, or apply Roundup spray solutions in containers or spray tanks made of galvanized or unlined steel (except stainless steel).
Asparagus Beans, edible Beet greens Beets, red Broccoli Cabbage Cauliflower Corn, sweet and pop Horseradish, as a weed and as a crop Jerusalem artichoke	Perennial grass control, applications outside the growing season				
	Roundup	2-3 lb.	(See remarks)	(See remarks)	Use for quackgrass or Johnsongrass control. Apply to quackgrass when 6 to 8 in. tall in fall or spring. Apply to Johnsongrass when at least 12 in. tall and actively growing. Do not till until 3 to 7 days after application. Does not provide residual weed control. Do not mix, store, or apply Roundup spray solutions in galvanized steel or unlined steel containers (except stainless steel). For control of volunteer horseradish, apply 3 to 4 lb. in mid-September. Field should have been disced 4 to 6 weeks prior to application. For this control practice, use spray coverage only.
	Kale Lentils Lettuce Mustard greens Okra Onions Peas Potato, Irish and sweet		Radishes Spinach		

**\* Restricted-use herbicide.**

NOTE: In the suggestions in this publication, trade names of herbicides are usually used. The list below shows trade names and their corresponding common names. **Restricted-use herbicides are identified with an asterisk(\*).**

Common name	Trade name	Common name	Trade name	Common name	Trade name
alachlor .....	Lasso	dinoseb .....	Premerge-3, Sinox, Dinoseb	metribuzin .....	Lexone, Sencor
atrazine.....	AAtrex and Atrazine	diphenamid .....	Enide	napropamide .....	Devrinol
benfen .....	Balan	diuron .....	Karmex and others	naptalam .....	Alanap L
bensulide .....	Prefar	EPTC .....	Eptam	paraquat*..	Paraquat,* Gramoxone*
bentazon .....	Basagran	EPTC+safener.....	Eradicane	pronamide* .....	Kerb*
bromoxynil .....	Brominal	EPTC + safener	+ extender .....	propachlor .....	Ramrod, Propachlor
butylate+safener .....	Sutan+			pyrazon .....	Pyramin
CDAA .....	Randex			oryzalin .....	Surflan
chloramben .....	Amibex	fluchloralin .....	Basalin	simazine .....	Princep
chlorpropham .....	Furloe	glyphosate .....	Roundup, Kleen Up	terbacil .....	Sinbar
cyanazine .....	Bladex	linuron .....	Lorox	trifluralin .....	Treflan
cycloate .....	Ro-Neet	MCPA, MCPB....	Vacate, numerous	Petroleum solvent..	Stoddard Solvent
dalapon .....	Dowpon	metolachlor .....	Dual	2,4-D (amine).....	(numerous)
DCPA .....	Dacthal				

### Storing Pesticides and Containers

Keep pesticides and containers in a separate building, room, or enclosure used only for this purpose. Such building or rooms should be dry, ventilated, and locked. Fence outside storage areas to protect children and animals and to discourage pilferage. CAUTION: Do not store weedkillers, herbicides, or defoliant in the same room with insecticides. Chlorate salts can create a fire or explosion hazard. Remove only the pesticides needed for one day's operation and return empty containers — and any unused pesticide — to the storage area each day.

### Disposing of Pesticides and Containers

**Surplus pesticides.** To dispose of surplus pesticide mixtures, try to find other areas with the same pest problem and use up any extra tank mix or rinse water on these areas. Do not drain surplus pesticides in any location where they can contaminate wells, streams, rivers, lakes, or ponds.

Operators of landfills meeting environmental safety standards can obtain supplemental permits to handle toxic waste materials, including pesticides. To dispose of large quantities of surplus pesticides, contact the Illinois EPA Division of Land Pollution Control to locate the nearest landfill with a supplemental permit for toxic waste or to obtain specific instructions about disposal.

**Pesticide containers.** All empty pesticide containers, regardless of their type, should be rinsed three times before disposal. Rinse water should be dumped in the tank. Triple-rinsed containers should be punctured or broken to facilitate drainage and to prevent reuse for any purpose. They should then be hauled to a sanitary landfill for disposal. Small quantities of containers may be buried singly in open fields, with due regard for the protection of surface and subsurface water.

Illinois regulations permit the burning of combustible containers provided that they are burned on the premises where they were used, that they are burned more than 1,000 feet from residential areas, that the burning will not cause undue visibility or environmental hazards, and that no reasonable alternative disposal method is available.

Do not breathe smoke from burning pesticide containers, and do not burn containers that have weedkillers such as 2,4-D or similar herbicides. When these change to a gas, the vapors may damage nearby crops and shrubbery. Pesticides containing chlorates may explode when heated and therefore should not be burned.

### Growers with Several Crops in a Small Area

Growers with several crops in a small area should be especially careful when applying herbicides. The tendency is to apply more if the quantity measured out "looks" as if it is not enough. A low-percentage granular formulation is suggested for small areas. *Check rates and application techniques on the container label very carefully. Applications must be accurate and uniform. Excessive amounts may cause injury to present or subsequent crops.*

Ideally, a specific herbicide should be fitted to a specific crop species. When growing several different crops in a small area, however, it is often impractical and expensive to use all the appropriate herbicides. Following are two herbicides that can be used on a wide range of vegetables.

**Dacthal.** Dacthal is cleared on a large number of vegetables. As listed in this circular, it may not always be the herbicide of preference. It can be used on broccoli, brussels sprouts, cauliflower, cabbage, snap beans, mung beans, Southern peas, soybeans, seeded melons, cucumbers, squash, collards, kale, mustard greens, turnips (root and greens), garlic, horseradish, onions, potatoes (Irish), sweet potatoes and yams, tomatoes, eggplant, peppers, and strawberries. *Do not use on beets or Swiss chard.*

Dacthal is a preemergence herbicide that must be applied to weed-free soil. It controls very small weed seedlings soon after the weed seeds germinate. It is most effective if rainfall occurs or if the soil is irrigated within 2 to 3 days after application.

A one-time application to all species is not always possible because some plants are susceptible to injury in early-growth stages. It is preferable to use Dacthal at seeding or transplanting time if the species is adapted to it. When this is not possible, the weeds should be removed and Dacthal applied to prevent further weed development. Consult the label for the appropriate application time.

Dacthal is effective in controlling annual grasses that are a problem in the spring. Broadleaf weeds that escape control should be mechanically removed.

**Treflan.** Treflan is widely available because it is used in soybean culture in Illinois. It can be purchased as a liquid with 4 pounds of active ingredient per gallon for large garden areas or as a low-percentage granular formulation for smaller areas. The amount of Treflan to use correlates very closely with the type of soil. The appropriate amounts are shown on the container label.

Treflan can be used for weed control in beans (green, lima, and dry), broccoli, brussels sprouts, cabbage, cauliflower, carrots, kale, mustard greens, okra, peas, peppers, tomatoes, and turnip greens. *Treflan may injure sweet corn.*

*Treflan must be mixed with or watered into the soil to prevent loss of the chemical from the soil surface.* A rototiller, disc, or similar implement should be used to mix Treflan with the soil to a 3- to 4-inch depth. When it is impractical to mix Treflan with the soil mechanically, remove all germinated and growing weeds and allow the herbicide to be carried into the soil through rainfall or sprinkle irrigation.

Treflan is quite effective on annual grasses, but many broadleaf weeds will need to be mechanically removed.

### **Mulches — see paragraph 3, page 1**

### **Other Publications on Weed Control**

Copies of the following publications on weed control are available from the office of your county Extension adviser in agriculture and the Office of Agricultural Publications, 123 Mumford Hall, 1301 West Gregory Drive, Urbana, Illinois 61801.

Calibrating and Adjusting Granular Row Applicators — Circular 1008

Controlling Weeds in the Home Garden — Circular 1051

Equipment and Calibration: Low Pressure Sprayers — Circular 1192

Turfgrass Pest Control — Circular 1076


Herbicides for Commercial Fruit Crops in Illinois — H-659

Herbicide Recommendations for Commercial Nurserymen — NC-2-80

1983 Row Crop Weed Control Guide


Weed Control in Small Grains, Forages, and Pastures







UNIVERSITY OF ILLINOIS-URBANA  
Q. 630.71L6C C005  
CIRCULAR URBANA, ILL.  
907 REV. 1983



3 0112 019541488