

HIGHLY ACCURATE QUANTUM-CHEMICAL CALCULATIONS FOR THE INTERSTELLAR MOLECULES C_3 AND $l-C_3H^+$

PETER BOTSCHWINA, BENJAMIN SCHRÖDER, CHRISTOPHER STEIN, PETER SEBALD, RAINER OSWALD, *Institute of Physical Chemistry, Georg-August-Universität Göttingen, Göttingen, Germany.*

Composite potential energy surfaces with coupled-cluster contributions up to CCSDTQP were constructed for C_3 and $l-C_3H^+$ and used in the calculation of spectroscopic properties. The use of very large AO basis sets and the consideration of higher-order correlation beyond CCSD(T) is of utmost importance for C_3 in order to arrive at quantitative spectroscopic data. The first detection of $l-C_3H^+$ in the interstellar medium was reported by Pety et al.,^a who attributed 9 radio lines observed in the horsehead photodissociation region to that species. That assignment was questioned by the recent theoretical work of Huang et al.^b However, our more accurate calculations are well in support of the original assignment. The calculated ground-state rotational constant is $B_0 = 11248$ MHz, only 0.03% off from the radio astronomical value of 11244.9512 ± 0.0015 MHz. The ratio of centrifugal distortion constants $D_0(\text{exp.})/D_e(\text{theor.})$ of 1.8 is quite large, but reasonable in comparison with C_3O^c and C_3 .

^aJ. Pety, P. Gratier, V. Guzmán, E. Roueff, M. Gerin et al., *Astron. Astrophys.* **2012**, A68, 1-8.

^bX. Huang, R. C. Fortenberry, T. J. Lee, *Astrophys. J. Lett.* **2013**, 768:L25, 1-5.

^cP. Botschwina, R. Oswald, *J. Chem. Phys.* **2008**, 129, 044305.