

CONSISTENT ASSIGNMENT OF THE VIBRATIONS OF MONOHALOSUBSTITUTED BENZENES

JOE HARRIS, ANNA ANDREJEVA, WILLIAM DUNCAN TUTTLE, *School of Chemistry, University of Nottingham, Nottingham, United Kingdom*; IGOR PUGLIESI, CHRISTIAN SCHRIEVER, *Fakultät für Physik, Ludwig-Maximilians-Universität, München, Deutschland*; TIM WRIGHT, *School of Chemistry, University of Nottingham, Nottingham, United Kingdom*.

When substituted benzenes become a focus of a spectroscopic study there are various well known vibrational labelling schemes present,^{a,b} however it was shown in recent works the description of monohalobenzene vibrations in terms of benzene modes (ie. Wilson notation) is questionable in some cases.^{c,d} A new scheme is presented which uses the motions of monofluorobenzene vibrations as a basis for labelling vibrational assignments of monosubstituted benzenes.^d The scheme has been successfully applied to the ground and excited states of toluene and its deuterated-methyl group isotopologue.^{e,f} Here we present the application of the scheme to fluorobenzene and its fully deuterated analogue. One-colour resonance-enhanced multiphoton ionization (REMPI) spectroscopy was employed in order to characterise the fluorobenzene and fluorobenzene-*d*₅ excited state.

^aE. B. Wilson Jr., *Phys. Rev.*, 45, 706 (1934)

^bG. Varsanyi, *Assignments of the Vibrational Spectra of Seven Hundred Benzene Derivatives*, Wiley, New York, 1974, Vol. I and II

^cI. Pugliesi, N. C. Tonge and M. C. R. Cockett, *J. Chem. Phys.*, 129, 104303 (2008)

^dA. M. Gardner and T. G. Wright, *J. Chem. Phys.*, 135, 114305 (2011)

^eA. M. Gardner, A. M. Green, V. M. Tame-Reyes, V. H. K. Wilton and T. G. Wright, 138, 134303 (2013)

^fA. M. Gardner, A. M. Green, V. M. Tame-Reyes, K. L. Reid, J. A. Davies, V. H. K. Wilton and T. G. Wright, manuscript accepted