

THE EFFECT OF TERMINAL SUBSTITUTION ON THE HELICAL CARBON STRUCTURE OF FLUORO-ALKANE CHAINS: A PURE ROTATIONAL STUDY OF  $\text{CH}_2\text{OH}-\text{C}_{n-1}\text{F}_{2n-1}$  ( $n = 4, 5, \& 6$ )

AARON Z. A. SCHWARTZ, *Natural and Social Science, Purchase College SUNY, Purchase, NY, USA*;  
MARK P. MATURO, DANIEL A. OBENCHAIN, *Department of Chemistry, Wesleyan University, Middletown, CT, USA*; S. A. COOKE, *Natural and Social Science, Purchase College SUNY, Purchase, NY, USA*.

Continuing a series of studies to investigate the change in structure of hydrocarbons as the amount of fluorination is increased to varying degrees of substitution, we present a survey on the change in the helical nature of the fluorinated carbon backbone when a  $-\text{CH}_2\text{OH}$  group is substituted for a terminal  $-\text{CF}_3$  group. Spectra for 1H,1H-heptafluorobutan-1-ol, 1H,1H-nonafluoropentan-1-ol, and 1H,1H-undecafluorohexan-1-ol were collected separately using a chirped-pulse FTMW spectrometer in the range of 7-13 GHz. Only one conformation was observed for each molecule. Additional measurements of the 1H,1H-heptafluorobutan-1-ol were completed using a Balle-Flygare cavity instrument. Assignments of the singly-substituted  $^{13}\text{C}$  isotopologues of the 1H,1H-heptafluorobutan-1-ol were also measured. A comparison of both *ab initio* and experimental structures will be presented.