

TOWARD ROTATIONAL STATE-SELECTIVE PHOTOIONIZATION OF ThF⁺ IONS

YAN ZHOU, KIA BOON NG, DAN GRESH, WILLIAM CAIRNCROSS, MATT GRAU, YIQI NI, ERIC CORNELL, JUN YE, *JILA, National Institute of Standards and Technology and Univ. of Colorado Department of Physics, University of Colorado, Boulder, Boulder, CO, USA.*

ThF⁺ has been chosen to replace HfF⁺ for a second-generation measurement of the electric dipole moment of the electron (eEDM). Compared to the currently running HfF⁺ eEDM experiment, ThF⁺ has several advantages: (i) the eEDM-sensitive state (³Δ₁) is the ground state, which facilitates a long coherence time [1]; (ii) its effective electric field (35 GV/cm) is 50% larger than that of HfF⁺, which promises a direct increase of the eEDM sensitivity [2]; and (iii) the ionization energy of neutral ThF is lower than its dissociation energy, which introduces greater flexibility in rotational state-selective photoionization via core-nonpenetrating Rydberg states [3]. In this talk, we first present our strategy of preparing and utilizing core-nonpenetrating Rydberg states for rotational state-selective ionization. Then, we report spectroscopic data of laser-induced fluorescence of neutral ThF, which provides critical information for multi-photon ionization spectroscopy.

[1] D. N. Gresh, K. C. Cossel, Y. Zhou, J. Ye, E. A. Cornell, *Journal of Molecular Spectroscopy*, 319 (2016), 1-9

[2] M. Denis, M. S. Nørby, H. J. A. Jensen, A. S. P. Gomes, M. K. Nayak, S. Knecht, T. Fleig, *New Journal of Physics*, 17 (2015) 043005.

[3] Z. J. Jakubek, R. W. Field, *Journal of Molecular Spectroscopy* 205 (2001) 197–220.