

OBSERVATION OF HEAVY RYDBERG STATES IN H₂ AND HD

MAXIMILIAN BEYER, FREDERIC MERKT, *Laboratorium für Physikalische Chemie, ETH Zurich, Zurich, Switzerland.*

The binding energies of the hydrogen atom are given by the Rydberg formula

$$E_n = -\frac{\mathcal{R}_\infty \mu / m_e}{(n - \delta)^2},$$

where the quantum defect δ vanishes in the case of a pure Coulomb potential.

Heavy Rydberg systems can be realized when the electron is replaced by an anion, which leads in the case of H⁺H⁻ to an almost 1000 times larger Rydberg constant and to an infinite number of vibrational states. In the diabatic molecular basis, these ion-pair states are described by long-range Coulomb potentials with ¹Σ_g⁺ and ¹Σ_u⁺ symmetry. In this basis, the level energies are described by an almost energy-independent, nonzero quantum defect, reflecting the finite size of H⁻. Strong interactions at small internuclear distances lead to strong variation of δ with n .

Gerade [2] and ungerade [3] ion-pair states have been observed in H₂ with principal quantum numbers up to $n = 240$. The quantum defects in this range were found to vary with energy, indicating the inadequacy of a pure diabatic picture.

Spectra of ungerade heavy Rydberg states of H₂ with $n = 160 - 520$ showing that the quantum defect only becomes energy independent for $n > 350$ will be presented, supporting the description using a diabatic basis.

I will also present first observations of ion-pair states in HD, showing two series of heavy Rydberg states, H⁺D⁻ and H⁻D⁺, which have different series limits. The experimental results will be discussed and compared with calculations using both an adiabatic and a diabatic basis.

[1] S. Pan, and F. H. Mies, *J. Chem. Phys.* **89**, 3096 (1988).

[2] M. O. Vieitez, T. I. Ivanov, E. Reinhold, C. A. de Lange, and W. Ubachs, *Phys. Rev. Lett.* **101**, 163001 (2008).

[3] R. C. Ekey, and E. F. McCormack, *Phys. Rev. A* **84**, 020501(R) (2011).