

A CANONICAL APPROACH TO GENERATE MULTIDIMENSIONAL POTENTIAL ENERGY SURFACES

JAY R. WALTON, *Department of Mathematics, Texas A & M University, College Station, TX, USA*;
LUIS A. RIVERA-RIVERA, *Department of Physical Sciences, Ferris State University, Big Rapids, MI, USA*;
ROBERT R. LUCCHESI, *Department of Chemistry, Texas A & M University, College Station, TX, USA*.

Previously adaptations of canonical approaches were applied to algebraic forms of the classic Morse, Lennard-Jones, and Kratzer potentials. Using the classic Morse, Lennard-Jones, or Kratzer potential as reference, inverse canonical transformations allow the accurate generation of Born-Oppenheimer potentials for H_2^+ ion, neutral covalently bound H_2 , van der Waals bound Ar_2 , and the hydrogen bonded 1-dimensional dissociative coordinate in water dimer. This methodology is now extending to multidimensional potential energy surfaces, and as a proof-of-concept, it is applied to the 3-dimensional water molecule potential surface. Canonical transformations previously developed for diatomic molecules are used to construct accurate approximations to the 3-dimensional potential surface of the water molecule from judiciously chosen 1-dimensional planar slices that are shown to have the same canonical shape as the classical Lennard-Jones potential curve. Spline interpolation is then used to piece together the 1-dimensional canonical potential curves, to obtain the full 3-dimensional potential surface of water molecule with a relative error less than 0.008.