

## TRACING THE ORIGINS OF NITROGEN BEARING ORGANICS TOWARD ORION KL WITH ALMA

BRANDON CARROLL, *Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA, USA*; NATHAN CROCKETT, *Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA*; OLIVIA H. WILKINS, *Chemistry, California Institute of Technology, Pasadena, CA, USA*; EDWIN BERGIN, *Department of Astronomy, University of Michigan, Ann Arbor, MI, USA*; GEOFFREY BLAKE, *Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA, USA*.

A comprehensive analysis of a broadband 1.2 THz wide spectral survey of the Orion Kleinmann-Low nebula (Orion KL) has shown that nitrogen bearing complex organics trace systematically hotter gas than O-bearing organics toward this source. The origin of this O/N dichotomy remains a mystery. If complex molecules originate from grain surfaces, N-bearing species may be more difficult to remove from grain surfaces than O-bearing organics. Theoretical studies, however, have shown that hot ( $T=300$  K) gas phase chemistry can produce high abundances of N-bearing organics while suppressing the formation of O-bearing complex molecules. In order to distinguish these distinct formation pathways we have obtained extremely high angular resolution observations of methyl cyanide ( $\text{CH}_3\text{CN}$ ) using the Atacama Large Millimeter/Submillimeter Array (ALMA) toward Orion KL. By simultaneously imaging  $^{13}\text{CH}_3\text{CN}$  and  $\text{CH}_2\text{DCN}$  we map the temperature structure and D/H ratio of  $\text{CH}_3\text{CN}$ . We will present updated results of these observations and discuss their implications for the formation of N-bearing organics in the interstellar medium.