

MICROWAVE CHARACTERIZATION OF THIOBENZALDEHYDE AND OTHER PRODUCTS IN THE DISCHARGE OF BENZENE WITH SULFUR ADDITIVES

VALENTINA DELL'ISOLA, *Dipartimento di Chimica G. Ciamician, Università di Bologna, Bologna, Italy*; JESSIE P PORTERFIELD, KELVIN LEE, *Radio and Geoastronomy Division, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA*; BRANDON CARROLL, *Atomic and Molecular Physics, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA*; CRISTINA PUZZARINI, *Dep. Chemistry 'Giacomo Ciamician', University of Bologna, Bologna, Italy*; MICHAEL C MCCARTHY, *Atomic and Molecular Physics, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA*.

Although many organo-sulfur species have been detected in molecular clouds and star-forming regions, they are insufficient to account for the expected sulfur content. In an attempt to identify new potential sulfur reservoirs, we have studied the discharge of benzene with both H₂S and CS₂. Thiobenzaldehyde is the first species we have characterized at high resolution using chirped pulse and cavity enhanced microwave spectroscopy in the 2-40 GHz region. Production of thiobenzaldehyde was particularly prominent in the discharge of benzene with carbon disulfide. Isotopic substitution with ¹³CS₂ indicates a relatively simple formation pathway for thiobenzaldehyde, initiated by attack of the benzene ring by ¹³CS followed by an H-atom shift from the benzene ring. Although much weaker, observation of thiobenzaldehyde in the discharge of benzene with H₂S suggests that other pathways may be relevant.