


PROPANE ISOTOPOLOGUES: HIGH RESOLUTION SYNCHROTRON FAR-IR SPECTRA OF THE SYMMETRICALLY DEUTERATED SPECIES  $\text{CH}_3\text{CH}_2\text{CD}_3$ ,  $\text{CD}_3\text{CH}_2\text{CD}_3$  AND  $\text{C}_3\text{D}_8$ . FIRST EXPERIMENTALLY DETERMINED GROUND STATE CONSTANTS FOR THESE SPECIES

STEPHEN J. DAUNT, ROBERT GRZYWACZ, *Department of Physics & Astronomy, The University of Tennessee-Knoxville, Knoxville, TN, USA*; COLIN WESTERN, *School of Chemistry, University of Bristol, Bristol, United Kingdom*; BRANT E. BILLINGHURST, JIANBAO ZHAO, *EFD, Canadian Light Source Inc., Saskatoon, Saskatchewan, Canada*.



We are continuing our project of obtaining high resolution vibration-rotation spectra of propane isotopologues using the Far-IR beamline at the Canadian National Synchrotron (CLS). We have already reported on all the singly  $^{13}\text{C}$  and singly D substituted varieties as well as the 2,2- $\text{D}_2$  species at previous ISMS meetings (2017-19) and in a recent paper on 2- $^{13}\text{C}$ -Propane<sup>a</sup>. These studies have allowed us to determine ground state inertial and centrifugal distortion rotational constants for these molecules that have no pure rotational spectra in the literature except for the 6 lines reported by Lide<sup>b</sup> in 1960 of normal propane and two  $^{13}\text{C}$  and D versions. In this talk we will present the high resolution spectra ( $0.00096\text{ cm}^{-1}$ ) for the CCC skeletal bendings of  $\text{CH}_3\text{CH}_2\text{CD}_3$ ,  $\text{CD}_3\text{CH}_2\text{CD}_3$  and  $\text{C}_3\text{D}_8$ . These studies have yielded corrected observed band origins<sup>c</sup> and rotational constants for the three species. Preliminary values for each species are listed here in wavenumbers.  $\text{CH}_3\text{CH}_2\text{CD}_3$ :  $\nu_9 = 335.664740(40)$ ,  $A_0 = 0.8185513(12)$ ,  $B_0 = 0.24400666(39)$ ,  $C_0 = 0.21852642(41)$ ;  $\text{CD}_3\text{CH}_2\text{CD}_3$ :  $\nu_{16} = 306.4$ ,  $A_0 = 0.711202$ ,  $B_0 = 0.213021$ ,  $C_0 = 0.193244$ ;  $\text{C}_3\text{D}_8$ :  $\nu_9 = 303.936065(23)$ ,  $A_0 = 0.58742224(42)$ ,  $B_0 = 0.20872437(23)$ ,  $C_0 = 0.18588200(18)$ .

<sup>a</sup>Daunt, Grzywacz, Western, Lafferty, Flaud, Billinghurst, and Hutchings, *J. Mol. Structure*, in press (doi:10.1016/j.molstruc.2020.127851).

<sup>b</sup>D. R. Lide, *J. Chem. Physics* 33, 1514-1518 (1960).

<sup>c</sup>Gayles and King, *Spectrochim. Acta* 21, 543-557 (1965); K. M. Gough, W. F. Murphy and K. Raghavachari, *J. Chem. Phys.* 87, 3332-3340 (1987).