

STRUCTURE OF THE MODEL GRIGNARD-TYPE REAGENT ClZnCH_3 ($\tilde{X}^1\text{A}_1$) BY MILLIMETER-WAVE SPECTROSCOPY

MARK BURTON, *Department of Chemistry and Biochemistry, University of Arizona, Tucson, AZ, USA;*
NAZIFA TABASSUM, *Department of Molecular and Cellular Biology, University of Arizona, Tucson, AZ, USA;*
LUCY M. ZIURYS, *Department of Chemistry and Biochemistry; Department of Astronomy, Arizona Radio Observatory, University of Arizona, Tucson, AZ, USA.*

Pure rotational spectra of the $^{37}\text{ClZnCH}_3$, ClZnCD_3 , and $\text{ClZn}^{13}\text{CH}_3$ isotopologues of monomeric ClZnCH_3 ($\tilde{X}^1\text{A}_1$) have been recorded using millimeter-wave direct absorption techniques in the frequency range 263 – 303 GHz. These species were synthesized in the gas phase in a DC discharge by the reaction of zinc vapor, produced in a Broida-type oven, with $^{37}\text{ClCH}_3$ (in natural chlorine abundance), ClCD_3 , or $\text{Cl}^{13}\text{CH}_3$. The data for each isotopologue were analyzed with a symmetric top Hamiltonian and rotational and centrifugal constants determined. In combination with previous measurements of $\text{Cl}^{64}\text{ZnCH}_3$, $\text{Cl}^{66}\text{ZnCH}_3$, and $\text{Cl}^{68}\text{ZnCH}_3$, an $r_m^{(2)}$ structure was determined for this organozinc compound. The bond lengths were calculated to be $r_{\text{Cl-Zn}} = 2.0831(1) \text{ \AA}$, $r_{\text{Zn-C}} = 1.9085(1) \text{ \AA}$, and $r_{\text{C-H}} = 1.1806(5) \text{ \AA}$. The H-C-H bond angle was found to be 110.5° – slightly larger than that in methane. These data serve to benchmark future structure calculations of organozinc compounds, which are widely used in organic synthesis.