

Radial segregation patterns in rotating granular mixtures: Waviness selection*

K. M. Hill, G. Gioia, and D. Amaravadi

*Department of Theoretical and Applied Mechanics,
University of Illinois, Urbana, IL 61801*

(Dated: December 13, 2004)

Abstract

We model the radial segregation patterns that form in a thin rotating drum partially filled with beads of two sizes. We predict that the waviness (or amplitude-to-wavelength ratio, denoted w) of a pattern should be subjected to low-pass filtering with a cut-off waviness w_c that depends strongly on the fill level of the drum. Then we perform experiments and find that $w \leq w_c$ for all patterns, in accord with our prediction. We also find that $w = w_c$ for (and only for) steady patterns, and conclude that the waviness of a steady pattern is selected by the low-pass filtering.

* To appear in *Physical Review Letters*

When a granular mixture is vibrated or made to flow, it segregates by type of grain [1]. For instance, if a bed of grains of different sizes is sheared parallel to the top surface, the larger grains segregate to the surface of the bed and the smaller ones to the bottom [2]. In nature, this form of granular segregation might explain why the gravelly bed of a piedmont stream is frequently topped with a self-organized, stabilizing *armour layer* composed of the largest rocks in the bed [3]. Granular segregation also occurs in numerous industrial processes, often with undesirable effects [4]. To study granular segregation, some researchers have used a thin rotating drum partially filled with beads of different sizes. Within the first rotation of the drum, a radial segregation pattern forms (Fig. 1a-c) [5]. The pattern of Fig. 1c forms for all fill levels of the drum; we call it a *moon pattern* because it reminds us of an ashy moon on a dark sky. For most fill levels, the moon pattern appears to be steady [6]. Yet *for fill levels close to 1/2* the moon pattern evolves through a series of transient, increasingly *wavy*, striped segregation patterns or *sun patterns* as the rotation continues (Fig. 1d-g) [7]. Upon attaining a value that depends strongly on the fill level, the *waviness* ceases to increase, and a sun pattern of the selected waviness remains steady thereafter (Fig. 1h) [6, 7]. Here we develop a model of waviness selection in sun patterns. In developing our model, we are led to identify the waviness of a sun pattern (as yet a loose visual descriptor) with the amplitude-to-wavelength ratio of its stripes. We find that in contrast to many systems in which a wavelength is selected at the onset of a runaway process (often embodied by an eigenproblem), in sun patterns the waviness is selected when a low-pass filter arrests a runaway process of waviness amplification. To test our findings, we perform experiments and compare the results with the predictions of the model.

In our experiments we use an acrylic drum (of radius $R = 15$ cm and thickness 8 mm) filled with water [8] and, up to a distance $R+d$ measured from the bottom of the drum, with a mixture of glass beads (Jaygo Inc., NJ; $\rho = 2.54$ g/cm³) (Fig. 2a). The average composition of the mixture is 40% 3 mm black beads and 60% 1 mm white beads, by volume (measured by weight). We rotate the drum about its axis with an angular velocity $\Omega = 1$ rpm and take pictures at regular intervals (e.g., Fig. 1). By processing these pictures [9], we obtain an index of the waviness, p^2/A , where p is the perimeter of the segregation pattern and A is the area contained within the perimeter. Fig. 2b shows a plot of p^2/A vs. the number of rotations for a fill level close to 1/2; within a few tens of rotations, p^2/A attains a steady value that depends strongly on the fill level (Fig. 2c).




FIG. 1: (a) The drum partially filled with well-mixed black and white beads. (b) The drum after 1/4 rotation, (c) 1/2 rotation (moon pattern), (d) 1 rotation, (e) 2 rotations, (f) 3 rotations, (g) 4 rotations, and (h) many rotations (steady sun pattern). In these experiments the free surface remains flat and steady as the drum rotates.




FIG. 2: (a) The drum filled to a distance $R + d$ measured from the bottom of the drum. The ratio $-1 \leq d/R \leq 1$ characterizes the fill level of the drum; if $d/R = 0$ the fill level is 1/2. (b) p^2/A vs. the number of rotations of the drum for a fill level close to 1/2. (c) The steady value of p^2/A vs. d/R for a range of fill levels. The steady value of p^2/A peaks for a fill level slightly higher than 1/2 ($d/R = 0.1$). For a moon pattern at a fill level of 1/2, $p^2/A = 2(2 + \pi)^2/\pi \approx 17$.




FIG. 3: (a) The flowing layer of maximum depth b is bounded by the free surface PQ and by the arc PMQ. The point O represents the axis of the drum. (b) A partially formed moon pattern. (c) A moon pattern. The value of x depends on the fill level, the average composition of the mixture, and b . (d) If $d > b$, the moon pattern includes a circle (with center at the axis of the drum) containing mixed beads that never enter the flowing layer. Cf. the pictures of Fig. 2c.

To shed light on these results, we start by describing how the beads move in the rotating drum. The beads move relative to one another only in a shallow surficial layer of maximum depth b known as the *flowing layer* (Fig. 3a) [10]. Below the flowing layer the beads—and, with the beads, the segregation pattern—move in solidlike rotation with the drum. The flowing layer exchanges beads with the stratum below through the arc PMQ of Fig. 3a. The beads positioned along the arc PM can be said to be *thawing* as they enter the flowing layer and cease to move in solidlike rotation with the drum. Thus we call PM the *thawing arc* and the flux of beads that enters the flowing layer through PM the *thawing flux*. Likewise, we call MQ the *freezing arc* and the flux of beads that leaves the flowing layer through MQ the *freezing flux*. Consider now the experiment of Fig. 1a-c. Since the black and the white beads are initially well mixed, they enter the flowing layer in a thawing flux of constant composition (i.e., a thawing flux whose average composition is constant in time). As they move in the flowing layer the larger (black) beads segregate to the upper part of the layer, and the smaller beads to the lower part. When these beads leave the flowing layer, they begin to form a moon pattern (Fig. 3b); when they reenter the flowing layer (again in a thawing flux of constant composition), the moon pattern becomes complete (Fig. 3c-d). This argument [5] explains the formation of moon patterns for all fill levels. For future reference, we stress that *a thawing flux of constant composition leads to the rapid formation of a moon pattern*.

Before turning to the sun patterns, we simplify the geometry of the thawing and freezing arcs. We substitute the thawing arc PM with the straight thawing line LM and the freezing arc MQ with the straight freezing line MR (Fig. 4a). The thawing and freezing lines are




FIG. 4: (a) LM=the thawing line, MR=the freezing line. Here M is below the axis of the drum ($d' < 0$). Beads that freeze at the same time remain on a straight line (b) as they rotate with the drum (c). (d) Idealized sun pattern. The value of x pertains to the moon pattern (Fig 3c). This figure shows the sequence in which the beads leave and reenter the flowing layer for $d' < 0$. If we turn the page up side down, this same figure will show the sequence in which the beads leave and reenter the flowing layer for $d' > 0$ (but with the thawing line on the right and the freezing line on the left). Because the geometry is the same regardless of the sign of d' , the equations that we derive based on this geometry apply to all fill levels. (e) A bead that freezes at a distance $x - \xi$ to the right of M thaws at distance $x - \xi$ to the left of M. Suppose that two beads freeze simultaneously, the 1st at $\xi = 0$ and the 2nd at $\xi = \xi'$; if the 1st thaws at a time t , the 2nd thaws at a time $t - \tau(\xi')$, where $\tau(\xi) = 2\xi \sin \theta / v$. (f) Plots of r vs. $|\psi|$ for c_f equal to one term (curve 'A') and 20 terms ('B') of the Fourier expansion of $S_T(t)$. 'C' is an idealization of 'B.' A high w gives a thawing flux of constant composition (g), a low w a thawing flux of oscillating composition (h).

parallel to the free surface at a distance $|d'|$ from the axis of the drum, where $d' = d - b$. The beads positioned along the freezing line MR at a given time remain on a straight line while they rotate with the drum (Fig. 4b-c).

Consider now the idealized sun pattern of Fig. 4d. For a time interval $T/2$, the beads

that leave the flowing layer through the segment Δx of the freezing line are black; then, for a time interval $T/2$, these beads are white, and so on in a cyclic fashion. Thus the beads leave the flowing layer in a freezing flux of oscillating composition. The stripes unfold in the fanlike arrangement of Fig. 4c; they have an amplitude Δx and a wavelength $\lambda = vT$, where $v = \Omega x \sec \theta$ (Fig. 4d). Fig. 4e illustrates the sequence in which the beads of Fig. 4d leave (and later reenter) the flowing layer.

With the help of Fig. 4e, we seek a mathematical characterization of the thawing and freezing fluxes. We start by defining the local composition $-1 \leq c \leq 1$ of a mixture in the form $c = f_b - f_w$, where f_b and f_w is the local volume fraction of black and white beads, respectively. On the freezing (or thawing) line, the local composition c may be a function of the position ξ and the time t , $c = c(\xi, t)$. (For the idealized sun pattern of Fig. 4, on the freezing line $c = c_f(\xi, t) = S_T(t)$, where $S_T(t)$ is an oscillating square function of amplitude 2 and period T , and on the thawing line $c = c_t(\xi, t) = S_T(t - \tau(\xi))$, where $\tau(\xi) = 2\xi \sin \theta / v$ is a time shift, see the caption to Fig. 4e.) The flux of c advected through the segment Δx of the freezing (or thawing) line is $\varphi(t) = v_n \int_{-\Delta x/2}^{\Delta x/2} c(\xi, t) d\xi$, where $v_n = \Omega x$ is the component of v normal to the freezing (or thawing) line [11]. The mean value of $\varphi(t)$, $\langle \varphi \rangle = \frac{1}{T} \int_0^T \varphi(t) dt$, is zero, because in the idealized sun pattern the white stripes have the same volume as the black stripes [12]. On the other hand, the root mean square of $\varphi(t)$, $\sqrt{\langle \varphi^2 \rangle} = \sqrt{\frac{1}{T} \int_0^T \varphi^2(t) dt}$, is zero for a flux of constant composition, but positive for a flux of oscillating composition, and provides a measure of the amplitude of the compositional oscillations of the flux $\varphi(t)$. Thus, if $\varphi_f(t)$ is the freezing flux and $\varphi_t(t)$ is the thawing flux, $r = \sqrt{\langle \varphi_t^2 \rangle / \langle \varphi_f^2 \rangle}$ measures the extent to which the compositional oscillations of the freezing flux carry over to the thawing flux. (For example, $r < 1$ indicates that the the fluxes becomes less oscillating with each freezing-and-thawing cycle.) To compute r , we take $c_f(\xi, t) = 4 \cos(2\pi t / T) / \pi$ (the first term in the Fourier expansion of $S_T(t)$), and therefore $c_t(\xi, t) = 4 \cos(2\pi(t - \tau(\xi)) / T) / \pi$, where $\tau(\xi) = 2\xi \sin \theta / v$. The result is $r = \sin |\psi| / |\psi|$, where $\psi = 2\pi w \sin \theta$ and $w = \Delta x / \lambda$, i.e., the amplitude-to-wavelength ratio or *waviness*. Curve ‘A’ in Fig. 4f is a plot of r vs. $|\psi|$. We have also computed r by taking c_f equal to the first twenty terms in the Fourier expansion of $S_T(t)$, with a similar result (curve ‘B’ in Fig. 4f). In both curve ‘A’ and curve ‘B’ r equals 1 and is *stationary* for $|\psi| = 0$, and then r decreases for increasing values of $|\psi|$. Thus, except for small values of $|\psi|$, for which $r \approx 1$, $r < 1$ and the flux tends to become less oscillating with each freezing-and-thawing cycle. For simplicity, we choose a ψ_c and assume $r = 1$ for

$|\psi| < \psi_c$ and $r = 0$ for $|\psi| > \psi_c$ (curve ‘C’ in Fig. 4f). With the reasonable choice $\psi_c = \pi/5$, this translates into $r = 1$ for $w < w_c$ and $r = 0$ for $w > w_c$, where $w_c = |\csc \theta|/10$ or

$$w_c = \frac{1}{10} \sqrt{1 + \left(\frac{x}{d-b} \right)^2}. \quad (1)$$

For $w > w_c$, $r = 0$ and the beads enter the flowing layer in a thawing flux of constant composition (to see this point, it is helpful to study a case with very high waviness, e.g. Fig 4g), leading to the rapid formation of a *moon* pattern. Thus, a sun pattern of waviness $w > w_c$ is wiped out. On the other hand, for $w < w_c$, $r = 1$ and the beads enter the flowing layer in a thawing flux of oscillating composition (to see this point, it is helpful to study a case with very low waviness, e.g. Fig 4h). Thus, a sun pattern of waviness $w < w_c$ may persist. We conclude that the waviness of a sun pattern is subjected to low-pass filtering with the cut-off waviness w_c given by (1). The cut-off waviness depends on the fill level, the average composition of the mixture, and the depth of the flowing layer through the variables d , x , and b . The peak value of w_c occurs for $d = b$ (or $d' = 0$). This is consistent with the experiments of Fig. 2c, in which w peaks for $d = 0.1R$; in fact, we have measured the depth of the flowing layer to be $b \approx 0.1R$ in those experiments.

To test these predictions, we compute $d' = d - b$ (using $b = 0.1R$), d'/x , and w [13] for each of the patterns, transient or steady, that we photographed in the experiments of Fig. 2c. Then, we plot the point $(w, d'/x)$ for each pattern (Fig. 5) and the curve w_c vs. d'/x predicted by the model. The effect of the low-pass filtering is manifest in Fig. 5: the experimental points lie beneath the theoretical curve. Further, for any given value of d'/x the waviness draws near the cut-off waviness as the pattern becomes steady [14], indicating that the waviness of a steady pattern is *selected* by the low-pass filtering. To understand how this comes about, we turn our attention back to Fig. 2c, in which a moon pattern has just formed. Now this moon pattern is likely to be slightly wavy; if the drum continues to rotate, the initially low waviness will be amplified (Fig. 2b). (The simulations of Khakhar *et al.* [7] indicate that the amplification takes place *inside* the flowing layer.) By virtue of this amplification, the waviness will increase up to the cut-off value w_c —whereupon the filtering will equilibrate the amplification, bringing the amplification to an end [15]. (Note that for most fill levels w_c is very low, so that the amplification will be arrested shortly after the formation of the moon pattern, which will appear to be steady.)

To summarize: As soon as a moon pattern forms, it is subjected to waviness amplifica-




FIG. 5: A comparison of the predictions of the model with the experimental results. The curve represents the predicted cut-off waviness. Each point represents an experimental measurement. Points with the same $(d - b)/x$ correspond to the same experiment; successive points with the same $(d - b)/x$ approach the cut-off waviness as the pattern in the experiment becomes steady.

tion and low-pass filtering. In time, the filtering arrests the amplification, thereby selecting the waviness of the steady sun pattern. Here we have developed a model of waviness filtering, ignoring the amplification. We have been able to do so because the filtering and the amplification are spatially disjoint: the filtering stems from the way in which the beads move *outside* the flowing layer, whereas the amplification stems from the way in which the beads move *inside* the flowing layer [7]. Thus the filtering depends only on the geometry of the patterns and remains independent of the segregation mechanisms that operate in the

flowing layer. On the other hand, the amplification may depend on the difference in bead sizes, the density contrast between the beads and the interstitial fluid, and other factors that pertain to the segregation mechanisms that operate in the flowing layer. We shall take up the development of a model of waviness amplification in a separate paper.

We are grateful for the financial support of the Research Board and the CRI Program at UIUC.

-
- [1] A. Rosato, K. J. Strandburg, F. Prinz, and R. H. Swendsen, *Phys. Rev. Lett.* **58**, 1038 (1987); J. B. Knight, H. M. Jaeger, and S. R. Nagel, *ibid.* **70**, 3728 (1993); O. Zik, D. Levine, S. G. Shtrikman, and J. Stavans, *ibid.* **73**, 644 (1994); S. B. Savage and C. K. K. Lun, *J. Fluid Mech.* **189**, 311 (1988); H. A. Makse, S. Havlin, P. R. King, and H. E. Stanley, *Nature (London)* **386**, 379 (1997).
 - [2] This might be the first form of granular segregation ever noted in a scientific paper; A. Schoklitsch, *Akad. Wiss. (Vienna)* **142**, 343 (1933).
 - [3] The armour layer has often been ascribed to selective erosion (e.g., R. Bettess and A. Frangipane, *J. Hydr. Res.* **41**, 179 (2003)), but field data pointing to the importance of granular segregation has been known for a long time (e.g., L. B. Leopold, M. G. Wolman and J. P. Miller, *Fluvial Processes in Geomorphology*, ch. 7 (W. H. Freeman and Co., San Francisco, 1964)).
 - [4] T. Shinbrot and F. J. Muzzio, *Phys. Today* **53**, No. 3, 25 (2000).
 - [5] E. Clement, J. Rajchenbach, J. Duran, *Europhys. Lett.* **30**, 7 (1995); F. Cantelaube and D. Bideau, *Europhys. Lett.* **30**, 133 (1995); D. V. Khakhar, J. J. McCarthy, and J. M. Ottino, *Phys. Fluids* **9**, 3600 (1997).
 - [6] K. M. Hill *et al.*, *PNAS* **96**, 11701 (1999).
 - [7] D. V. Khakhar, A. V. Orpe, and J. M. Ottino, *Powder Tech.* **116**, 232 (2001).
 - [8] If air is the interstitial fluid instead of water, similar patterns form, and the sun patterns occur only in a narrow range of fill levels [6, 7], just as in Fig. 2c. This fact suggests that our analysis could be broadly applicable, in particular to granular flows in air.
 - [9] We use NIH Image, which computes the length of the boundary between dark and light areas in a given picture. To calibrate this computation, we use the picture of a moon pattern, for

which p can be calculated analytically.

- [10] N. Jain, J. M. Ottino, and R. M. Lueptow, Phys. Fluids **14**, 572 (2002); D. Bonamy, F. Daviaud, and L. Laurent, *ibid.* **14**, 1666 (2002); K. M. Hill, G. Gioia, and V. V. Tota, Phys. Rev. Lett. **91**, 064302 (2003).
- [11] We take v_n to be constant on the segment Δx of the freezing (or thawing) line.
- [12] In our model, a moon pattern becomes wavy when the interface between the black and the white beads, initially a circular arc, becomes undulated with a wavelength λ . Thus the volume of the white stripes is the same as the volume of the black stripes, regardless of the average composition of the mixture.
- [13] To estimate w we write $p = 2x + L + 2n\Delta x$, where n is the number of stripes, and $L = (\pi + 2\theta)x \sec \theta$ (Fig. 4d). By substituting $n = 2L/\lambda$ we arrive to the formula $w = (p - 2x - L)/2L$. From the picture of the pattern we obtain p , x , and L , and then estimate w using this formula. For a moon pattern $p = 2x + L$, and $w = 0$.
- [14] The case $d'/x = 0$ (which corresponds to $|\psi| = 0$) is an exception, because w_c is singular for $d'/x = 0$. Nevertheless, there exists an upper bound, w_u , on the attainable waviness, and we can *regularize* the cut-off waviness by taking $w_c = w_u$ for $d'/x = 0$. To obtain an expression for w_u , we note that for $d' = 0$ (or $d = b$) the depth of the flowing layer is d , and its half-length R . Thus we can estimate $\Delta x \sim R$ and $\lambda \sim 2b = 2d$, and therefore $w_u \sim R/2d$; for the experiments of Fig. 2c, $d/R = 0.1$ and we have $w_u \sim 5$, which is in reasonable accord with the peak value of w measured in those experiments (Fig. 5).
- [15] We can write an equation of equilibrium between the filtering and the amplification in the form $r_a r = 1$, where $0 < r < 1$ represents the filtering and $r_a > 1$ represents the amplification. For $w = w_c$ (or $\psi = \psi_c$), r can take any value between 0 and 1 (Fig. 4f), and the filtering can equilibrate any value of $r_a > 1$.

List of Recent TAM Reports

No.	Authors	Title	Date
973	Pushkin, D. O., and H. Aref	Self-similarity theory of stationary coagulation – <i>Physics of Fluids</i> 14 , 694–703 (2002)	July 2001
974	Lian, L., and N. R. Sottos	Stress effects in ferroelectric thin films – <i>Journal of the Mechanics and Physics of Solids</i> (submitted)	Aug. 2001
975	Fried, E., and R. E. Todres	Prediction of disclinations in nematic elastomers – <i>Proceedings of the National Academy of Sciences</i> 98 , 14773–14777 (2001)	Aug. 2001
976	Fried, E., and V. A. Korchagin	Striping of nematic elastomers – <i>International Journal of Solids and Structures</i> 39 , 3451–3467 (2002)	Aug. 2001
977	Riahi, D. N.	On nonlinear convection in mushy layers: Part I. Oscillatory modes of convection – <i>Journal of Fluid Mechanics</i> 467 , 331–359 (2002)	Sept. 2001
978	Sofronis, P., I. M. Robertson, Y. Liang, D. F. Teter, and N. Aravas	Recent advances in the study of hydrogen embrittlement at the University of Illinois – Invited paper, Hydrogen–Corrosion Deformation Interactions (Sept. 16–21, 2001, Jackson Lake Lodge, Wyo.)	Sept. 2001
979	Fried, E., M. E. Gurtin, and K. Hutter	A void-based description of compaction and segregation in flowing granular materials – <i>Continuum Mechanics and Thermodynamics</i> , in press (2003)	Sept. 2001
980	Adrian, R. J., S. Balachandar, and Z.-C. Liu	Spanwise growth of vortex structure in wall turbulence – <i>Korean Society of Mechanical Engineers International Journal</i> 15 , 1741–1749 (2001)	Sept. 2001
981	Adrian, R. J.	Information and the study of turbulence and complex flow – <i>Japanese Society of Mechanical Engineers Journal B</i> , in press (2002)	Oct. 2001
982	Adrian, R. J., and Z.-C. Liu	Observation of vortex packets in direct numerical simulation of fully turbulent channel flow – <i>Journal of Visualization</i> , in press (2002)	Oct. 2001
983	Fried, E., and R. E. Todres	Disclinated states in nematic elastomers – <i>Journal of the Mechanics and Physics of Solids</i> 50 , 2691–2716 (2002)	Oct. 2001
984	Stewart, D. S.	Towards the miniaturization of explosive technology – Proceedings of the 23rd International Conference on Shock Waves (2001)	Oct. 2001
985	Kasimov, A. R., and Stewart, D. S.	Spinning instability of gaseous detonations – <i>Journal of Fluid Mechanics</i> (submitted)	Oct. 2001
986	Brown, E. N., N. R. Sottos, and S. R. White	Fracture testing of a self-healing polymer composite – <i>Experimental Mechanics</i> (submitted)	Nov. 2001
987	Phillips, W. R. C.	Langmuir circulations – <i>Surface Waves</i> (J. C. R. Hunt and S. Sajjadi, eds.), in press (2002)	Nov. 2001
988	Gioia, G., and F. A. Bombardelli	Scaling and similarity in rough channel flows – <i>Physical Review Letters</i> 88 , 014501 (2002)	Nov. 2001
989	Riahi, D. N.	On stationary and oscillatory modes of flow instabilities in a rotating porous layer during alloy solidification – <i>Journal of Porous Media</i> 6 , 1–11 (2003)	Nov. 2001
990	Okhuysen, B. S., and D. N. Riahi	Effect of Coriolis force on instabilities of liquid and mushy regions during alloy solidification – <i>Physics of Fluids</i> (submitted)	Dec. 2001
991	Christensen, K. T., and R. J. Adrian	Measurement of instantaneous Eulerian acceleration fields by particle-image accelerometry: Method and accuracy – <i>Experimental Fluids</i> (submitted)	Dec. 2001
992	Liu, M., and K. J. Hsia	Interfacial cracks between piezoelectric and elastic materials under in-plane electric loading – <i>Journal of the Mechanics and Physics of Solids</i> 51 , 921–944 (2003)	Dec. 2001
993	Panat, R. P., S. Zhang, and K. J. Hsia	Bond coat surface rumpling in thermal barrier coatings – <i>Acta Materialia</i> 51 , 239–249 (2003)	Jan. 2002
994	Aref, H.	A transformation of the point vortex equations – <i>Physics of Fluids</i> 14 , 2395–2401 (2002)	Jan. 2002
995	Saif, M. T. A, S. Zhang, A. Haque, and K. J. Hsia	Effect of native Al_2O_3 on the elastic response of nanoscale aluminum films – <i>Acta Materialia</i> 50 , 2779–2786 (2002)	Jan. 2002

List of Recent TAM Reports (cont'd)

No.	Authors	Title	Date
996	Fried, E., and M. E. Gurtin	A nonequilibrium theory of epitaxial growth that accounts for surface stress and surface diffusion — <i>Journal of the Mechanics and Physics of Solids</i> 51 , 487-517 (2003)	Jan. 2002
997	Aref, H.	The development of chaotic advection — <i>Physics of Fluids</i> 14 , 1315-1325 (2002); see also <i>Virtual Journal of Nanoscale Science and Technology</i> , 11 March 2002	Jan. 2002
998	Christensen, K. T., and R. J. Adrian	The velocity and acceleration signatures of small-scale vortices in turbulent channel flow — <i>Journal of Turbulence</i> , in press (2002)	Jan. 2002
999	Riahi, D. N.	Flow instabilities in a horizontal dendrite layer rotating about an inclined axis — <i>Journal of Porous Media</i> , in press (2003)	Feb. 2002
1000	Kessler, M. R., and S. R. White	Cure kinetics of ring-opening metathesis polymerization of dicyclopentadiene — <i>Journal of Polymer Science A</i> 40 , 2373-2383 (2002)	Feb. 2002
1001	Dolbow, J. E., E. Fried, and A. Q. Shen	Point defects in nematic gels: The case for hedgehogs — <i>Archive for Rational Mechanics and Analysis</i> , in press (2004)	Feb. 2002
1002	Riahi, D. N.	Nonlinear steady convection in rotating mushy layers — <i>Journal of Fluid Mechanics</i> 485 , 279-306 (2003)	Mar. 2002
1003	Carlson, D. E., E. Fried, and S. Sellers	The totality of soft-states in a neo-classical nematic elastomer — <i>Journal of Elasticity</i> 69 , 169-180 (2003) with revised title	Mar. 2002
1004	Fried, E., and R. E. Todres	Normal-stress differences and the detection of disclinations in nematic elastomers — <i>Journal of Polymer Science B: Polymer Physics</i> 40 , 2098-2106 (2002)	June 2002
1005	Fried, E., and B. C. Roy	Gravity-induced segregation of cohesionless granular mixtures — <i>Lecture Notes in Mechanics</i> , in press (2002)	July 2002
1006	Tomkins, C. D., and R. J. Adrian	Spanwise structure and scale growth in turbulent boundary layers — <i>Journal of Fluid Mechanics</i> (submitted)	Aug. 2002
1007	Riahi, D. N.	On nonlinear convection in mushy layers: Part 2. Mixed oscillatory and stationary modes of convection — <i>Journal of Fluid Mechanics</i> 517 , 71-102 (2004)	Sept. 2002
1008	Aref, H., P. K. Newton, M. A. Stremler, T. Tokieda, and D. L. Vainchtein	Vortex crystals — <i>Advances in Applied Mathematics</i> 39 , in press (2002)	Oct. 2002
1009	Bagchi, P., and S. Balachandar	Effect of turbulence on the drag and lift of a particle — <i>Physics of Fluids</i> , in press (2003)	Oct. 2002
1010	Zhang, S., R. Panat, and K. J. Hsia	Influence of surface morphology on the adhesive strength of aluminum/epoxy interfaces — <i>Journal of Adhesion Science and Technology</i> 17 , 1685-1711 (2003)	Oct. 2002
1011	Carlson, D. E., E. Fried, and D. A. Tortorelli	On internal constraints in continuum mechanics — <i>Journal of Elasticity</i> 70 , 101-109 (2003)	Oct. 2002
1012	Boyland, P. L., M. A. Stremler, and H. Aref	Topological fluid mechanics of point vortex motions — <i>Physica D</i> 175 , 69-95 (2002)	Oct. 2002
1013	Bhattacharjee, P., and D. N. Riahi	Computational studies of the effect of rotation on convection during protein crystallization — <i>International Journal of Mathematical Sciences</i> , in press (2004)	Feb. 2003
1014	Brown, E. N., M. R. Kessler, N. R. Sottos, and S. R. White	<i>In situ</i> poly(urea-formaldehyde) microencapsulation of dicyclopentadiene — <i>Journal of Microencapsulation</i> (submitted)	Feb. 2003
1015	Brown, E. N., S. R. White, and N. R. Sottos	Microcapsule induced toughening in a self-healing polymer composite — <i>Journal of Materials Science</i> (submitted)	Feb. 2003
1016	Kuznetsov, I. R., and D. S. Stewart	Burning rate of energetic materials with thermal expansion — <i>Combustion and Flame</i> (submitted)	Mar. 2003

List of Recent TAM Reports (cont'd)

No.	Authors	Title	Date
1017	Dolbow, J., E. Fried, and H. Ji	Chemically induced swelling of hydrogels – <i>Journal of the Mechanics and Physics of Solids</i> , in press (2003)	Mar. 2003
1018	Costello, G. A.	Mechanics of wire rope – Mordica Lecture, Interwire 2003, Wire Association International, Atlanta, Georgia, May 12, 2003	Mar. 2003
1019	Wang, J., N. R. Sottos, and R. L. Weaver	Thin film adhesion measurement by laser induced stress waves – <i>Journal of the Mechanics and Physics of Solids</i> (submitted)	Apr. 2003
1020	Bhattacharjee, P., and D. N. Riahi	Effect of rotation on surface tension driven flow during protein crystallization – <i>Microgravity Science and Technology</i> 14 , 36–44 (2003)	Apr. 2003
1021	Fried, E.	The configurational and standard force balances are not always statements of a single law – <i>Proceedings of the Royal Society</i> (submitted)	Apr. 2003
1022	Panat, R. P., and K. J. Hsia	Experimental investigation of the bond coat rumpling instability under isothermal and cyclic thermal histories in thermal barrier systems – <i>Proceedings of the Royal Society of London A</i> 460 , 1957–1979 (2003)	May 2003
1023	Fried, E., and M. E. Gurtin	A unified treatment of evolving interfaces accounting for small deformations and atomic transport: grain-boundaries, phase transitions, epitaxy – <i>Advances in Applied Mechanics</i> 40 , 1–177 (2004)	May 2003
1024	Dong, F., D. N. Riahi, and A. T. Hsui	On similarity waves in compacting media – <i>Horizons in Physics</i> , in press (2003)	May 2003
1025	Liu, M., and K. J. Hsia	Locking of electric field induced non-180° domain switching and phase transition in ferroelectric materials upon cyclic electric fatigue – <i>Applied Physics Letters</i> 83 , 3978–3980 (2003)	May 2003
1026	Liu, M., K. J. Hsia, and M. Sardela Jr.	In situ X-ray diffraction study of electric field induced domain switching and phase transition in PZT-5H – <i>Journal of the American Ceramics Society</i> (submitted)	May 2003
1027	Riahi, D. N.	On flow of binary alloys during crystal growth – <i>Recent Research Development in Crystal Growth</i> , in press (2003)	May 2003
1028	Riahi, D. N.	On fluid dynamics during crystallization – <i>Recent Research Development in Fluid Dynamics</i> , in press (2003)	July 2003
1029	Fried, E., V. Korchagin, and R. E. Todres	Biaxial disclinated states in nematic elastomers – <i>Journal of Chemical Physics</i> 119 , 13170–13179 (2003)	July 2003
1030	Sharp, K. V., and R. J. Adrian	Transition from laminar to turbulent flow in liquid filled microtubes – <i>Physics of Fluids</i> (submitted)	July 2003
1031	Yoon, H. S., D. F. Hill, S. Balachandar, R. J. Adrian, and M. Y. Ha	Reynolds number scaling of flow in a Rushton turbine stirred tank: Part I – Mean flow, circular jet and tip vortex scaling – <i>Chemical Engineering Science</i> (submitted)	Aug. 2003
1032	Raju, R., S. Balachandar, D. F. Hill, and R. J. Adrian	Reynolds number scaling of flow in a Rushton turbine stirred tank: Part II – Eigen-decomposition of fluctuation – <i>Chemical Engineering Science</i> (submitted)	Aug. 2003
1033	Hill, K. M., G. Gioia, and V. V. Tota	Structure and kinematics in dense free-surface granular flow – <i>Physical Review Letters</i> , in press (2003)	Aug. 2003
1034	Fried, E., and S. Sellers	Free-energy density functions for nematic elastomers – <i>Journal of the Mechanics and Physics of Solids</i> 52 , 1671–1689 (2004)	Sept. 2003
1035	Kasimov, A. R., and D. S. Stewart	On the dynamics of self-sustained one-dimensional detonations: A numerical study in the shock-attached frame – <i>Physics of Fluids</i> (submitted)	Nov. 2003
1036	Fried, E., and B. C. Roy	Disclinations in a homogeneously deformed nematic elastomer – <i>Nature Materials</i> (submitted)	Nov. 2003
1037	Fried, E., and M. E. Gurtin	The unifying nature of the configurational force balance – <i>Mechanics of Material Forces</i> (P. Steinmann and G. A. Maugin, eds.), in press (2003)	Dec. 2003
1038	Panat, R., K. J. Hsia, and J. W. Oldham	Rumpling instability in thermal barrier systems under isothermal conditions in vacuum – <i>Philosophical Magazine</i> , in press (2004)	Dec. 2003

List of Recent TAM Reports (cont'd)

No.	Authors	Title	Date
1039	Cermelli, P., E. Fried, and M. E. Gurtin	Sharp-interface nematic-isotropic phase transitions without flow – <i>Archive for Rational Mechanics and Analysis</i> 174 , 151–178 (2004)	Dec. 2003
1040	Yoo, S., and D. S. Stewart	A hybrid level-set method in two and three dimensions for modeling detonation and combustion problems in complex geometries – <i>Combustion Theory and Modeling</i> (submitted)	Feb. 2004
1041	Dienberg, C. E., S. E. Ott-Monsivais, J. L. Ranchero, A. A. Rzeszutko, and C. L. Winter	Proceedings of the Fifth Annual Research Conference in Mechanics (April 2003), TAM Department, UIUC (E. N. Brown, ed.)	Feb. 2004
1042	Kasimov, A. R., and D. S. Stewart	Asymptotic theory of ignition and failure of self-sustained detonations – <i>Journal of Fluid Mechanics</i> (submitted)	Feb. 2004
1043	Kasimov, A. R., and D. S. Stewart	Theory of direct initiation of gaseous detonations and comparison with experiment – <i>Proceedings of the Combustion Institute</i> (submitted)	Mar. 2004
1044	Panat, R., K. J. Hsia, and D. G. Cahill	Evolution of surface waviness in thin films via volume and surface diffusion – <i>Journal of Applied Physics</i> (submitted)	Mar. 2004
1045	Riahi, D. N.	Steady and oscillatory flow in a mushy layer – <i>Current Topics in Crystal Growth Research</i> , in press (2004)	Mar. 2004
1046	Riahi, D. N.	Modeling flows in protein crystal growth – <i>Current Topics in Crystal Growth Research</i> , in press (2004)	Mar. 2004
1047	Bagchi, P., and S. Balachandar	Response of the wake of an isolated particle to isotropic turbulent cross-flow – <i>Journal of Fluid Mechanics</i> (submitted)	Mar. 2004
1048	Brown, E. N., S. R. White, and N. R. Sottos	Fatigue crack propagation in microcapsule toughened epoxy – <i>Journal of Materials Science</i> (submitted)	Apr. 2004
1049	Zeng, L., S. Balachandar, and P. Fischer	Wall-induced forces on a rigid sphere at finite Reynolds number – <i>Journal of Fluid Mechanics</i> (submitted)	May 2004
1050	Dolbow, J., E. Fried, and H. Ji	A numerical strategy for investigating the kinetic response of stimulus-responsive hydrogels – <i>Journal of the Mechanics and Physics of Solids</i> (submitted)	June 2004
1051	Riahi, D. N.	Effect of permeability on steady flow in a dendrite layer – <i>Journal of Porous Media</i> , in press (2004)	July 2004
1052	Cermelli, P., E. Fried, and M. E. Gurtin	Transport relations for surface integrals arising in the formulation of balance laws for evolving fluid interfaces – <i>Journal of Fluid Mechanics</i> (submitted)	Sept. 2004
1053	Stewart, D. S., and A. R. Kasimov	Theory of detonation with an embedded sonic locus – <i>SIAM Journal on Applied Mathematics</i> (submitted)	Oct. 2004
1054	Stewart, D. S., K. C. Tang, S. Yoo, M. Q. Brewster, and I. R. Kuznetsov	Multi-scale modeling of solid rocket motors: Time integration methods from computational aerodynamics applied to stable quasi-steady motor burning – <i>Proceedings of the 43rd AIAA Aerospace Sciences Meeting and Exhibit</i> (January 2005), Paper AIAA-2005-0357 (2005)	Oct. 2004
1055	Ji, H., H. Mourad, E. Fried, and J. Dolbow	Kinetics of thermally induced swelling of hydrogels – <i>International Journal of Solids and Structures</i> (submitted)	Dec. 2004
1056	Fulton, J. M., S. Hussain, J. H. Lai, M. E. Ly, S. A. McGough, G. M. Miller, R. Oats, L. A. Shipton, P. K. Shreeman, D. S. Widrevitz, and E. A. Zimmermann	Final reports: Mechanics of complex materials, Summer 2004 (K. M. Hill and J. W. Phillips, eds.)	Dec. 2004
1057	Hill, K. M., G. Gioia, and D. R. Amaravadi	Radial segregation patterns in rotating granular mixtures: Waviness selection – <i>Physical Review Letters</i> , in press (2004)	Dec. 2004