

## CO<sub>2</sub> DIMER: FOUR INTERMOLECULAR MODES OBSERVED VIA INFRARED COMBINATION BANDS

JALAL NOROOZ OLIAEE, *Department of Physics and Astronomy, University of Calgary, Calgary, AB, Canada*; MEHDI DEGHANY, *Department of Mathematics, Physics and Engineering, Mount Royal University, Calgary, Canada*; MOJTABA REZAEI, *Department of Physics and Astronomy, University of Calgary, Calgary, AB, Canada*; BOB McKELLAR, *Steacie Laboratory, National Research Council of Canada, Ottawa, ON, Canada*; NASSER MOAZZEN-AHMADI, *Physics and Astronomy/Institute for Quantum Science and Technology, University of Calgary, Calgary, AB, Canada*.

Study of the carbon dioxide dimer has a long history, but there is only one previous observation of an intermolecular vibration [1]. Here we analyze four new combination bands of (CO<sub>2</sub>)<sub>2</sub> in the CO<sub>2</sub>  $\nu_3$  region ( $\sim 2350$  cm<sup>-1</sup>), observed using tunable infrared lasers and a pulsed slit-jet supersonic expansion. The previous combination band at 2382.2 cm<sup>-1</sup> was simple to assign [1]. A much more complicated band ( $\sim 2370$  cm<sup>-1</sup>) turns out to involve *two* upper states, one at 2369.0 cm<sup>-1</sup> ( $B_u$  symmetry), and the other at 2370.0 cm<sup>-1</sup> ( $A_u$ ). The spectrum can be nicely fit by including the Coriolis interactions between these states. Another complicated band around 2443 cm<sup>-1</sup> also involves two nearby upper states which are highly perturbed in so-far unexplained ways (possibly related to tunneling shifts).

With the help of new *ab initio* calculations [2], we assign the results as follows. The 2369.0 cm<sup>-1</sup> band is the combination of the forbidden  $A_g$  intramolecular fundamental (probably [1] at about 2346.76 cm<sup>-1</sup>) and the intermolecular geared bend ( $B_u$ ). The 2370.0 cm<sup>-1</sup> band is the combination of the same  $A_g$  fundamental and the intermolecular torsion ( $A_u$ ). This gives about 22.3 and 23.2 cm<sup>-1</sup> for the geared bend and torsion. The previous 2382.2 cm<sup>-1</sup> band [1] is the allowed  $B_u$  fundamental (2350.771 cm<sup>-1</sup>) plus two quanta of the geared bend ( $B_u$ ), giving 31.509 cm<sup>-1</sup> for this overtone. The highly perturbed 2442.7 cm<sup>-1</sup> band is the  $B_u$  fundamental plus the antigeared bend ( $A_g$ ), giving about 91.9 cm<sup>-1</sup> for the antigeared bend. Finally, the perturbed 2442.1 cm<sup>-1</sup> band is due to an unknown combination of modes which gains intensity from the antigeared bend by a Fermi-type interaction. Calculated values [2] are: 20.64 (geared bend), 24.44 (torsion), 32.34 (geared bend overtone), and 92.30 cm<sup>-1</sup> (antigeared bend), in good agreement with experiment.

[1] M. Deghany, A.R.W. McKellar, Mahin Afshari, and N. Moazzen-Ahmadi, *Mol. Phys.* **108**, 2195 (2010).

[2] X.-G. Wang, T. Carrington, Jr., and R. Dawes, private communication.