

## FAR-INFRARED SPECTROSCOPY OF *SYN*-VINYL ALCOHOL

PAUL RASTON, *Department of Chemistry and Biochemistry, James Madison University, Harrisonburg, Virginia, USA*; HAYLEY BUNN, *School of Chemistry and Physics, The University of Adelaide, Adelaide, South Australia, Australia*.

Vinyl alcohol has been extensively studied in both the microwave<sup>a,b</sup> and mid-IR<sup>c,d</sup> spectral regions, where 9 out of 15 vibrational modes have been identified. Here we present the first far-IR spectrum of vinyl alcohol, collected below 700 cm<sup>-1</sup> at the Australian Synchrotron. The high resolution (0.001 cm<sup>-1</sup>) spectrum reveals the  $\nu_{11}$  and  $\nu_{15}$  fundamentals of *syn*-vinyl alcohol at 489 cm<sup>-1</sup> and 407 cm<sup>-1</sup>, in addition to two hot bands of the  $\nu_{15}$  mode at 369 cm<sup>-1</sup> and 323 cm<sup>-1</sup>. High  $J$  transitions in the R-branch of the  $\nu_{15}$  band were found to be perturbed by an  $a$ -axis Coriolis interaction with the nearby  $\nu_{11}$  state. The  $\nu_{15}$  torsional mode of *syn*-vinyl alcohol was fit using a Watson's A-reduced Hamiltonian to yield rotational, centrifugal distortion, and Coriolis coupling parameters.

---

<sup>a</sup>S. Saito, Chem. Phys. Lett. 42, 3 (1976)

<sup>b</sup>M. Rodler et al., J. Am. Chem. Soc. 106, 4029 (1948)

<sup>c</sup>Y. Koga et al., J. Mol. Spec. 145, 315 (1991)

<sup>d</sup>D-L. Joo et al., J. Mol. Spec. 197, 68 (1999)